

Climate Change and Variability - Horizons Region

Prepared for Horizons Regional Council

September 2016

Prepared by:

Petra Pearce
Vijay Paul
Brett Mullan
Christian Zammit
Abha Sood
Rob Bell
Cliff Law

For any information regarding this report please contact:

Petra Pearce
Climate Scientist
Climate Applications
+64-9-375 2052
petra.pearce@niwa.co.nz

National Institute of Water & Atmospheric Research Ltd
Private Bag 99940
Viaduct Harbour
Auckland 1010

Phone +64 9 375 2050

NIWA CLIENT REPORT No: AKL2016-011
Report date: September 2016
NIWA Project: ELF16204

Quality Assurance Statement		
	Reviewed by:	Andrew Tait – Principal Scientist, Wellington
	Formatting checked by:	Andrew Tait – Principal Scientist, Wellington
	Approved for release by:	Ken Becker – Regional Manager, Auckland

© All rights reserved. This publication may not be reproduced or copied in any form without the permission of the copyright owner(s). Such permission is only to be given in accordance with the terms of the client's contract with NIWA. This copyright extends to all forms of copying and any storage of material in any kind of information retrieval system.

Whilst NIWA has used all reasonable endeavours to ensure that the information contained in this document is accurate, NIWA does not give any express or implied warranty as to the completeness of the information contained herein, or that it will be suitable for any purpose(s) other than those specifically contemplated during the Project or agreed by NIWA and the Client.

Contents

Executive summary	7
1 Introduction	12
2 Background: Global Climate Change – Science and Impacts.....	14
2.1 The Physical Science Basis (IPCC Working Group I)	14
2.2 Impacts, Adaptation and Vulnerability (IPCC Working Group II).....	16
2.3 Mitigation of Climate Change (IPCC Working Group III).....	17
3 Background: New Zealand Climate Change – Science and Impacts.....	18
3.1 Sectoral Impacts	18
4 Present Climate of Horizons Region.....	21
4.1 Spatial Patterns in Horizons Region’s Climate	21
4.2 Temporal Variability in Horizon Region’s Climate	24
4.2.1 Temperature.....	24
4.2.2 Rainfall.....	25
4.2.3 Drought.....	25
4.2.4 Extreme Rainfall Events.....	28
4.2.5 Growing Degree Days.....	29
4.3 Natural factors causing fluctuation in climate patterns over New Zealand	29
4.3.1 The effect of El Niño and La Niña	30
4.3.2 The effect of the Interdecadal Pacific Oscillation.....	31
4.3.3 The effect of the Southern Annular Mode	33
4.4 New Zealand Sea Level Trends and Variability	34
5 Projections of Horizons Region’s Future Climate	36
5.1 Horizons Climate Change Temperature Projections.....	36
5.1.1 Projections of Growing Degree-Days	41
5.1.2 Water Temperature Trends.....	41
5.2 Projections for Frosts and Hot Days under Climate Change.....	42
5.3 Horizons Climate Change Precipitation Projections	46
5.4 Temperature and precipitation projection comparisons within RCPs	54
5.5 Scenarios for Changes in Extreme Rainfall	58
5.6 Evaporation, Soil Moisture and Drought	62
5.7 Pressure and Wind.....	66

5.8	Hill Country Erosion	68
5.8.1	Climate change impacts on sediment loads in rivers	70
5.9	Solar Radiation.....	70
5.10	Climate Change and Sea Level	71
5.10.1	Effect of Sea-Level Rise on High Tide Exceedance Frequency.....	73
5.11	Climate Change Impacts on Other Coastal Hazard Drivers	76
5.12	Hydrological Impacts of Climate Change	77
5.13	Ocean Acidification	79
5.13.1	New Zealand-specific Impacts of Ocean Acidification	82
5.14	Considering both Anthropogenic and Natural Changes	83
6	Horizons Region – Impacts, Vulnerability and Adaptation	85
7	Acknowledgements	87
8	References.....	88

Tables

Table 4-1:	Historical relative sea-level rise rates.	35
Table 5-1:	Projected changes in seasonal and annual mean temperature (in °C) for the Horizons Region for 2040 and 2090.	41
Table 5-2:	Projected changes in the number of hot days and cold nights (frosts) at 2040 and 2090 for Horizons Region.	45
Table 5-3:	Projected changes in seasonal and annual precipitation (in %) between 1986-2005 and 2031-2050.	50
Table 5-4:	Projected changes in seasonal and annual precipitation (in %) between 1986-2005 and 2081-2100.	51
Table 5-5:	Augmentation factors (percentage increases per degree of warming) used in deriving changes in extreme rainfall for preliminary scenario studies.	59
Table 5-6:	Current rainfall depth-duration-frequency statistics for Palmerston North from HIRDS V3.	59
Table 5-7:	Projected rainfall depth-duration-frequency statistics for Palmerston North in 2100, for a low-range temperature scenario (1°C warming), from HIRDS V3.	60
Table 5-8:	Projected rainfall depth-duration-frequency statistics for Palmerston North in 2100 for a mid-range temperature scenario (2°C warming), from HIRDS V3.	60
Table 5-9:	Projected rainfall depth-duration-frequency statistics for Palmerston North in 2100, for a higher-end temperature scenario (3°C warming), from HIRDS V3.	60
Table 5-10:	Data analysis of effect of sea-level rise at Foxton on components of tide.	75
Table 5-11:	Projected changes in flow variables of the Manawatu River (in %) between 1986-2005 and 2031-2050 (2040) and 2081-2100 (2090).	77

Figures

Figure 1-1:	The Horizons Regional Council area. The region administered by the Council is outlined.	13
Figure 2-1:	CMIP5 multi-model simulated time series from 1950-2100 for change in global annual mean surface temperature relative to 1986-2005.	15
Figure 4-1:	Annual average temperature for the Horizons region (median for 1981-2010). ©NIWA.	22
Figure 4-2:	Annual rainfall for the Horizons region (median for 1981-2010). ©NIWA.	22
Figure 4-3:	Homogenised annual temperature time series for Whanganui from 1938 to 2014.	24
Figure 4-4:	Annual rainfall total (mm) at Whanganui from 1890 to 2014 (top) and Palmerston North from 1944 to 2013 (bottom).	25
Figure 4-5:	PED accumulation from 1947-2015 (July-June years) for Taumarunui (agent number 2250).	26
Figure 4-6:	PED accumulation from 1939-2015 (July-June years) for Wanganui, Spriggins Park EWS (agent number 3715).	27
Figure 4-7:	PED accumulation from 1940-2015 (July-June years) for Palmerston N (agent number 3238) and Palmerston North EWS (agent number 21963) - combined series.	27
Figure 4-8:	PED accumulation from 1950-2015 (July-June years) for Dannevirke (agent number 2534).	28
Figure 4-9:	Number of days per year with >2mm, >5mm and >10mm rain at Whanganui, Spriggins Park EWS site.	28
Figure 4-10:	Median annual growing degree-days in Horizons Region, 1981-2010. Left: Base 5°C, Right: Base 10°C.	29
Figure 4-11:	Time series of the Southern Oscillation Index from 1930 to 2016. Blue shades are indicative of La Niña periods and red shades are indicative of El Niño periods.	30
Figure 4-12:	Average summer percentage of normal rainfall during El Niño (left) and La Niña (right).	31
Figure 4-13:	The Interdecadal Pacific Oscillation (IPO) index.	32
Figure 4-14:	Percentage change in average annual rainfall, for the 1978-1998 period compared to the 1960-1977 period.	32
Figure 4-15:	Time series of the Southern Annular Mode from transient experiments forced with time-varying ozone-depleting substances and greenhouse gases.	34
Figure 4-16:	Relative sea-level rise at four main New Zealand ports, 1900-2007. Modified after Hannah and Bell (2012).	35
Figure 5-1:	Projected seasonal temperature changes at 2040 (2031-2050 average).	37
Figure 5-2:	Projected seasonal temperature changes at 2090 (2081-2100 average).	38
Figure 5-3:	Projected seasonal temperature changes at 2040 (2031-2050 average).	39
Figure 5-4:	Projected seasonal temperature changes at 2090 (2081-2100 average).	40
Figure 5-5:	Projected increase in number of hot days per year ($T_{max} > 25^{\circ}\text{C}$) at 2040 & 2090 for RCP 4.5 (left panels) and RCP 8.5 (right panels), for Horizons Region.	43
Figure 5-6:	Projected decrease in number of cold nights per year ($T_{min} < 0^{\circ}\text{C}$) at 2040 and 2090 for RCP 4.5 (left) and RCP 8.5 (right), for Horizons Region.	44

Figure 5-7:	Projected seasonal precipitation changes (in %) at 2040 (2031-2050 average).	46
Figure 5-8:	Projected seasonal precipitation changes (in %) at 2090 (2081-2100 average).	47
Figure 5-9:	Projected seasonal precipitation changes (in %) at 2040 (2031-2050 average).	48
Figure 5-10:	Projected seasonal precipitation changes (in %) at 2090 (2081-2100 average).	49
Figure 5-11:	Projected seasonal temperature changes by 2040 (2031-2050) averaged over the Horizons Region, for the four RCPs.	54
Figure 5-12:	Projected seasonal temperature changes by 2090 (2081-2100) averaged over the Horizons Region, for the four RCPs.	55
Figure 5-13:	Projected seasonal rainfall changes by 2040 (2081-2100) for Whanganui, for the four RCPs.	56
Figure 5-14:	Projected seasonal rainfall changes by 2090 (2081-2100) for Whanganui, for the four RCPs.	56
Figure 5-15:	Projected seasonal rainfall changes by 2040 (2081-2100) for Dannevirke, for the four RCPs.	57
Figure 5-16:	Projected seasonal rainfall changes by 2090 (2081-2100) for Dannevirke for the four RCPs.	58
Figure 5-17:	Change in the magnitude of the 99th percentile of daily precipitation (in %).	62
Figure 5-18:	RCM-projected changes in Potential Evapotranspiration Deficit (PED, in mm accumulation over the July-June ‘water year’) for Horizons Region.	63
Figure 5-19:	RCM-projected changes in the annual number (in days) of “dry days” (precipitation below 1 mm/day).	66
Figure 5-20:	Change in the magnitude of the 99th percentile of daily-mean wind speed, for all four RCPs and three future time periods.	68
Figure 5-21:	Projections of global mean sea level rise over the 21st century relative to 1986-2005.	71
Figure 5-22:	AR5 global mean sea-level projections extended to 2120 spliced with annual MSL from Wellington and Auckland.	72
Figure 5-23:	The frequency of occurrence of high tides exceeding different present day tide marks at Foxton.	75
Figure 5-24:	Change in ocean surface pH.	80
Figure 5-25:	Projected mean surface pH for the NZ EEZ open ocean from a suite of six CMIP5 models, using the RCP 8.5 scenario.	81
Figure 5-26:	Time series of Surface seawater pCO ₂ (μ atm, top panel), pH (middle panel), and saturation state of the carbonate mineral, aragonite (lower panel).	81
Figure 5-27:	New Zealand Temperature - historical record and an illustrative schematic projection illustrating future year-to-year variability.	84
Figure 6-1:	Time scales and adaptation. Planning for human-induced climate change becomes increasingly important as one moves right along this line.	85

Executive summary

This report describes changes which may occur over the coming century in the climate of the region administered by the Horizons Regional Council, and outlines some possible impacts of these changes.

To set the context, we summarise key findings of the recent (2013-2014) global climate change assessment undertaken by the Intergovernmental Panel on Climate Change.

- Warming of the climate system is ‘unequivocal’, and most of the observed increase in globally averaged temperatures since the mid-20th century is very likely due to the increase in greenhouse gas concentrations caused by human activities.
- The IPCC updates projections for global and regional changes in temperature, sea level, and precipitation for the coming century, and points to an expected increase in the frequency of heavy rainfall events.
- Recent global warming is already having physical and biological effects in many parts of the world.
- Work assessed by the IPCC indicates that limiting future global warming to targets which are currently being discussed internationally would require substantial reductions in global greenhouse gas emissions from human activities.
- Continued emissions of greenhouse gases will cause further warming and changes in all parts of the climate system. There are four scenarios named RCPs (Representative Concentration Pathways) by the IPCC. These RCPs represent different climate change mitigation scenarios – one (RCP 2.6) leading to a very low emissions level (requiring removal of CO₂ from the atmosphere), two stabilisation scenarios (RCPs 4.5 and 6.0), and one (RCP 8.5) with very high greenhouse gas concentrations. Therefore, the RCPs represent a range of 21st century climate policies.

Next, information is summarised about expected New Zealand national and regional impacts of climate change, from the IPCC chapter on Australia and New Zealand.

- New Zealand has warmed by $0.09 \pm 0.03^\circ\text{C}$ per decade since 1909, with more heat waves, fewer frosts, more rain in the south and west of New Zealand, less rain in the north and east of the North and South Islands, and a rise in sea level since 1900 of $1.7 \pm 0.1 \text{ mm/yr}$.
- Ongoing vulnerability in New Zealand to extreme events is demonstrated by substantial economic losses caused by droughts, floods, fire, tropical cyclones, and hail. During the 21st century, New Zealand’s climate is virtually certain to warm further, with noticeable changes in extreme events.
- Heat waves and fire risk are virtually certain to increase in intensity and frequency. Floods, landslides, droughts, and storm surges are likely to become more frequent and intense, and snow and frost to become less frequent.

- Precipitation changes are projected to lead to increased runoff in the west and south of the South Island and reduced runoff in the northeast of the South Island, and the east and north of the North Island.
- The potential impacts of climate change on industry are likely to be substantial. New Zealand's predominantly hydroelectric power generation is vulnerable to precipitation variability, changes in snow cover are likely to have a significant impact on the ski industry, and pasture production may be impacted by warming and elevated CO₂.

Horizons Region's present climate is then described.

- An upward trend in mean temperature, consistent with the overall New Zealand warming through the 20th century, is apparent at the long-term climate monitoring site at Whanganui. There are substantial year to year fluctuations in temperature superimposed on this long-term trend, with some years being over 1.5°C different from others.
- There is also substantial year to year variation in rainfall. Whanganui exhibits annual rainfall totals ranging from around 600 mm up to more than 1200 mm, and Palmerston North rainfall varies from about 700 to 1100mm.
- Three natural fluctuations leading to year-to-year variations are the El Niño-Southern Oscillation (ENSO), the Interdecadal Pacific Oscillation (IPO), and the Southern Annular Mode (SAM). These factors also lead to fluctuations in sea level.

Projections for Horizons Region's future climate are then covered:

- Future temperature scenarios for 2040 (2031-2050 relative to 1986-2005) show that annual average temperatures across the region are projected to increase by between 0.7°C (RCP 2.6) and 1.1°C (RCP 8.5). By 2090 (2081-2100 relative to 1986-2005) annual average temperatures are projected to increase by between 0.7°C (RCP 2.6) and 3.1°C (RCP 8.5). The greatest warming is projected for summer or autumn (depending on the RCP) and the least warming is projected for spring. A slight acceleration in warming is projected for the second 50 years of the 21st century compared to the first 50 years under the higher emission scenarios, and inland areas tend to warm more. There are only modest spatial gradients in the warming relative to the mean increase. By 2090 for RCP 4.5, the greatest warming is expected to occur in the northern half of Horizons Region in autumn, and in winter in most of the region apart from the Central Plateau. For RCP 8.5 by 2090, most warming is expected for the part of the region west of the Ruahine and Tararua Ranges, and inland from the coast, in summer.
- Future precipitation projections indicate slightly more rainfall in spring and winter for much of the region west of the Ruahine and Tararua Ranges to 2040, and less rainfall in autumn and summer. By 2090 for RCP 4.5, more rainfall is projected for the western part of the region in spring, summer, and especially winter. By 2090 under RCP 8.5, the north-western

part of Horizons Region is projected to receive more rainfall (by up to 20%) in winter, and the eastern part of the region may receive less rainfall (by up to 20%) in winter.

- For many of the chosen locations in Horizons Region, there is no clear precipitation signal, even at 2090 under RCP 8.5. The ensemble-average is often less than $\pm 5\%$, with the model range (the 5th and 95th percentile values) varying between quite large (>10%) increases and decreases. By 2040 (2031-2050, relative to 1986-2005), winter is the season with the most precipitation change, with a small increase in the ensemble-average (up to 8% across the different locations and RCPs). Akitio and Owahanga are the only locations to show a negative precipitation change by 2040.
- By 2090, there is a clearer precipitation signal at most of the chosen locations. For most places, winter is still the season with the most precipitation change, with some locations projecting increases in the ensemble-average at around 15% (Ohakune has the highest projected winter precipitation increase of 18% at RCP 8.5). The direction of projected precipitation change is not uniform across the region; precipitation for Akitio and Owahanga is projected to decrease by ~15% by 2090 under RCP 8.5.
- Projections for Horizons Region for the coming century also include a substantial decrease in cold nights, an increase in the number of hot days, and an increase in the frequency of very heavy rainfall.
- For engineering purposes some scenarios for changes in rainfall depth/duration/frequency statistics are provided for Palmerston North.
- Potential Evapotranspiration Deficit (PED) is calculated for the region, and analysis is broken down into sub-regions. PED, in units of mm, can be thought of as the amount of rainfall needed in order to keep pastures growing at optimum levels. A larger increase in PED over time equates to more drought risk in the future. In Horizons Region, central and eastern parts of the region experience the largest increases in PED by 2040 and 2090 under both RCP 4.5 and RCP 8.5.
- An increase in drought frequency is projected for Horizons Region of about 5% for 2030-2050 and 10% for 2070-2090, compared to 1980-1999 levels. The Central Plateau is less likely to be affected as the remainder of the region by climate change-induced drought. These projections were calculated from the IPCC Fourth Assessment Report emissions scenarios and will be updated in due course.
- The frequency of extreme winds (99th percentile) over the 21st century is likely to increase in the lower North Island (including Horizons Region).
- Changes in annual rainfall patterns as well as extreme rainfall will impact rates of hillslope erosion and river sedimentation. The northern and western portions of Horizons Region are more susceptible to erosion than to the east of the Ruahine and Tararua Ranges. Climate change will reduce the long-term effectiveness of SLUI (Sustainable Land Use Initiative) if other management scenarios are not adopted.

- New guidance on planning for sea level is expected in due course from the Ministry for the Environment (the last update was published in 2008). In the interim we suggest using a minimum sea-level rise scenario of 0.5 m by the 2090s (2090 to 2099) relative to the 1980-1999 average for coastal planning, plus an assessment of sensitivity to possible higher mean sea levels. For longer-term considerations an allowance for further sea-level rise of 10 mm/year beyond 2100 is recommended.
- Currently, 10% of high tides at Foxton exceed the MHWS-10 level (Mean High Water Spring level that 10% of high tides exceed, excluding weather and climate effects). With 0.4 m of sea-level rise, approximately 60% of high tides will exceed this level and with 0.8 m of sea-level rise about 99% of high tides will exceed this level, again excluding weather and climate effects such as storm surge.
- The following hydrological projections for the Manawatu River are presented, derived from the national surface water hydrological model TopNet:
 - Negligible change in mean annual flow is expected to occur under all RCPs at both 2040 and 2090.
 - Mean annual flood is expected to increase across most RCPs at both time periods, which is consistent with (small) mean annual precipitation increases for locations near the Manawatu River.
 - Mean annual low flow (Q10) is expected to decrease across all RCPs at both time periods, except RCP 2.6 at 2090 (median = 1% increase). However, the range of model results is quite large. The largest median ensemble change is projected for 2090 under RCP 8.5, which is a reduction in mean annual low flow of 12%.
 - For average seasonal flows, the median model results for most RCPs at both time periods are within $\pm 5\%$, and the ranges of model projections are quite large. However, under RCP 8.5 at 2090 for summer, flows are projected to decrease by 14% (median model result). This is consistent with small projected decreases in summer precipitation for the area around the Manawatu River for the same RCP and time period. Spring flows are projected to slightly increase by 2040 but decrease by 2090 for most RCPs.
 - FRE3 is the average number of high flow events (floods) per year that exceed three times the median flow. Small decreases in the median model result for FRE3 are projected for all RCPs except for RCP 8.5 at 2040 (small increase), and large increases in FRE3 ($>10\%$) are projected for all RCPs except RCP 2.6 (no change) at 2090.
- The pH of the oceans around New Zealand is projected to decrease, consistent with global trends. The variability and rate of change in pH will differ in coastal waters as these are also

influenced by terrestrial factors and run-off. Changes in ocean pH may have significant impacts on New Zealand fisheries and aquaculture into the future.

The Council is referred to material published by the Ministry for the Environment for guidance on assessing likely vulnerability and impacts for Horizons Region of these projected climate changes, and for considering adaptation options. Relevant issues could include:

- Implications of sea-level rise and coastal change for planning and development in coastal areas.
- Implications of river flow and sedimentation changes, as well as changing flood regimes for river engineering planning.
- Implications of potential changes in rainfall and of drought frequency for water demand, availability and allocation (including planning for irrigation schemes and storage).
- Implications of projected changes in extreme rainfall, erosion risk, and coastal hazards for council roading and stormwater drainage infrastructure, lifelines planning, and civil defence and emergency management.
- Opportunities which climate change may bring for new horticultural crops – and infrastructure and land-use issues that might arise.
- Implications of climate change (including potential changes in flood frequency, extreme rainfall (influencing hillslope and riverbank erosion) and in coastal hazards) for land-use planning (e.g. erosion control measures).
- Implications for natural ecosystems and their management, both terrestrial and marine. This is especially relevant given the two National Parks and a number of forest parks in the region.
- Building consideration of climate change impacts and adaptation into council planning as outlined in MfE guidance. Also important is consultation and discussion with stakeholders (e.g. groups of farmers, iwi) to help them identify climate-related risks and ways of building resilience.

This Envirolink Medium Advice Grant did not allow any ‘new’ research to be undertaken. However, during the compilation of this report, gaps were identified where further research for Horizons Region would be useful in order to understand potential climate change-related impacts:

- Region-specific modelling of climate-induced sea-level rise and coastal hazard drivers such as storm surge and waves.
- River flow projections for additional rivers in Horizons Region.

1 Introduction

Horizons Regional Council applied for and received funding from the Envirolink Fund (Ministry of Business, Innovation, and Employment) for NIWA to undertake a review of climate change projections and potential impacts for the Horizons Region, since the publication of the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report in 2013 and 2014.

This report describes climate changes which may occur over the coming century for the region administered by the Horizons Regional Council, and outlines some possible impacts of these changes. The report does not address the issue of mitigation (reducing greenhouse gas emissions, or increasing “sinks” such as areas of growing forest), apart from a brief summary of recent findings of the IPCC.

Consideration is given to both natural variations in the climate and to changes which may result from increasing global concentrations of greenhouse gases caused by human activities. Climatic factors discussed include temperature, rainfall, wind, evaporation, and soil moisture. River flow variables are also considered.

Possible changes along the coast in sea level are also considered. Figure 1-1 shows the Horizons Regional Council area of administration.

Preparation of the report has been supported through an Envirolink medium advice grant. This did not fund any new data analysis, but enabled us to draw on information which is already available from various sources. Much of this information is very new, resulting from the latest assessments of the Intergovernmental Panel on Climate Change (IPCC, 2013, IPCC, 2014a, IPCC, 2014b), and scenarios for New Zealand generated by NIWA scientists based on downscaling from global climate model runs undertaken for these IPCC assessments (undertaken through NIWA’s core-funded Regional Modelling Programme). The climate change information presented in this report is entirely consistent with recently-updated climate change guidance produced for the Ministry of the Environment (Mullan et al., 2016).




Figure 1-1: The Horizons Regional Council area. The region administered by the Council is outlined.

2 Background: Global Climate Change – Science and Impacts

This section summarises some key findings from the 2013 and 2014 IPCC Fifth Assessment Reports (AR5) as contextual information for the discussion of past and future climate changes in Horizons Region to follow in this report.

2.1 The Physical Science Basis (IPCC Working Group I)

The Summary for Policymakers of the IPCC AR5 Working Group I Report (IPCC, 2013) emphasises the following points regarding changes to the climate system:

- Warming of the climate system is ‘unequivocal’, and since the 1950s, many of the observed climate changes are unprecedented over short and long timescales (decades to millennia). These changes include warming of the atmosphere and ocean, diminishing of ice and snow, sea-level rise, and increases in the concentration of greenhouse gases.
- The atmospheric concentrations of carbon dioxide, methane, and nitrous oxide have increased to levels unprecedented in at least the last 800,000 years. Carbon dioxide concentrations have increased by 40% since pre-industrial times, primarily from fossil fuel emissions and secondarily from net land use change emissions. The ocean has absorbed about 30% of the emitted anthropogenic carbon dioxide, causing ocean acidification.
- Climate change is already influencing the intensity and frequency of many extreme weather and climate events globally.
- It is extremely likely that human influence has been the dominant cause of the observed warming since the mid-20th century.

Continued emissions of greenhouse gases will cause further warming and changes in all parts of the climate system. There are four scenarios named RCPs (Representative Concentration Pathways) by the IPCC. These RCPs represent different climate change mitigation scenarios – one (RCP 2.6) leading to a very low emissions level (requiring removal of CO₂ from the atmosphere), two stabilisation scenarios (RCPs 4.5 and 6.0), and one (RCP 8.5) with very high greenhouse gas concentrations. Therefore, the RCPs represent a range of 21st century climate policies.

By the middle of the 21st century, the magnitudes of the projected climate changes are substantially affected by the choice of scenario. **Global surface temperature change for the end of the 21st century is likely to exceed 1.5°C relative to 1850–1900** for all scenarios except for the lowest emissions scenario (RCP 2.6).

In contrast to the Fourth IPCC Assessment Report which concentrated on projections for the end of the 21st century, the Fifth Assessment Report projects climate changes for earlier in the 21st century as well in its Summary for Policymakers. As such, **the global mean surface temperature change for the period 2016–2035 (relative to 1986–2005) will likely be in the range of 0.3 to 0.7°C**. This assumes that there will be no major volcanic eruptions (which may cause global cooling) and that total solar irradiance remains similar. Temperature increases are expected to be larger in the tropics and subtropics than in the southern mid-latitudes (i.e. New Zealand).

The full range of projected globally averaged temperature increases for all scenarios for 2081-2100 (relative to 1986-2005) is 0.3 to 4.8°C (Figure 2-1). As global temperatures increase, it is virtually certain that there will be more hot and fewer cold temperature extremes over most land areas. It is very likely that heat waves will occur with a higher frequency and duration. Furthermore, in general, the contrast in precipitation between wet and dry regions and wet and dry seasons will increase. With increases in global mean temperature, mid-latitude and wet tropical regions will experience more intense and more frequent extreme precipitation events by the end of the 21st century.




Figure 2-1: CMIP5 multi-model simulated time series from 1950-2100 for change in global annual mean surface temperature relative to 1986-2005. Time series of projections and a measure of uncertainty (shading) are shown for scenarios RCP 2.6 (blue) and RCP 8.5 (red). Black (grey shading) is the modelled historical evolution using historical reconstructed forcings. The mean and associated uncertainties averaged over 2081–2100 are given for all RCP scenarios as coloured vertical bars. The numbers of CMIP5 models used to calculate the multi-model mean is indicated. After IPCC (2013).

The global ocean will continue to warm during the 21st century. Eventually, heat will penetrate into the deep ocean and affect ocean circulation. Sea ice is projected to shrink and thin in the Arctic. Some scenarios project that late summer Arctic sea ice extent could almost completely disappear by the end of the 21st century, and a nearly ice-free Arctic Ocean in late summer before mid-century is likely under the most extreme scenario. Northern Hemisphere spring snow cover will decrease as global mean surface temperature increases. The global glacier volume (excluding glaciers on the periphery of Antarctica) is projected to decrease by 15–85% by the end of the 21st century under different scenarios.

Global mean sea level will continue to rise during the 21st century. All scenarios project that the rate of sea level rise will very likely exceed that observed during 1971–2010 due to increased ocean warming and higher loss of mass from glaciers and ice sheets. For all scenarios, **the total range of projected sea level rise for 2081-2100 (relative to 1986-2005) is 0.26-0.82 m.** It is virtually certain that global mean sea level rise will continue beyond 2100, with sea level rise due to thermal expansion expected to continue for many centuries. The range for mean sea level rise beyond 2100 for different scenarios is from less than 1 m to more than 3 m, but sustained mass loss by ice sheets would cause larger sea level rise. Sustained warming greater than a critical threshold could lead to the near complete loss of the Greenland ice sheet over a millennium or more, causing a global mean sea level rise of up to 7 m. Current estimates place this threshold between 1 and 4°C global mean warming with respect to pre-industrial mean temperatures.

Cumulative CO₂ emissions largely determine global mean surface warming by the late 21st century and further into the future. Even if emissions are stopped, most aspects of global climate change will persist for many centuries.

2.2 Impacts, Adaptation and Vulnerability (IPCC Working Group II)

The IPCC AR5 Working Group II Summary for Policymakers (IPCC, 2014a) concludes that in recent decades, changes in climate have caused impacts on natural and human systems on all continents and across the oceans. Specifically, these include impacts to hydrological systems with regards to snow and ice melt, changing precipitation patterns and resulting river flow and drought, as well as terrestrial and marine ecosystems, the incidence of wildfire, food production, livelihoods, and economies.

Changes in precipitation and melting snow and ice are altering hydrological systems and are driving changes to water resources in terms of quantity and quality. The flow-on effects from this include impacts to agricultural systems, in particular crop yields, which have experienced more negative impacts than positive due to recent climate change. In response to changes in climate, many species have shifted their geographical ranges, migration patterns, and abundances. Some unique and threatened systems, including ecosystems and cultures, are already at risk from climate change. With increased warming around 1°C, the number of such systems at risk of severe consequences is higher, and many species with limited adaptive capacity (e.g. coral reefs and Arctic sea ice) are subject to very high risks with additional warming of 2°C. In addition, climate change-related risks from extreme events, such as heat waves, extreme precipitation, and coastal flooding, are already moderate/high with 1°C additional warming. Risks associated with some types of extreme events (e.g. heat waves) increase further with higher temperatures.

There is also the risk of physical systems or ecosystems undergoing abrupt and irreversible changes under increased warming. At present, warm-water coral reef and Arctic ecosystems are showing warning signs of irreversible regime shifts. With additional warming of 1-2°C, risks increase disproportionately and become high under additional warming of 3°C due to the threat of global sea level rise from ice sheet loss.

Global climate change risks are significant with global mean temperature increase of 4°C or more above pre-industrial levels and include severe and widespread impacts on unique or threatened systems, substantial species extinction, large risks to global and regional food security, and the combination of high temperature and humidity compromising normal human activities, including growing food or working outdoors in some areas for parts of the year.

Impacts of climate change vary regionally, and impacts are exacerbated by uneven development processes. Marginalised people are especially vulnerable to climate change and also to some adaptation and mitigation responses. This has been observed during recent climate-related extremes, such as heat waves, droughts, floods, cyclones, and wildfires, where different ecosystems and human systems are significantly vulnerable and exposed to climate variability. In addition, aggregate economic damages accelerate with increasing temperature.

In many regions, climate change adaptation experience is accumulating across the public and private sector and within communities. Adaptation is becoming embedded in governmental planning and development processes, but at this stage there has been only limited implementation of responses to climate change.

The overall risks of climate change impacts can be reduced by limiting the rate and magnitude of climate change.

2.3 Mitigation of Climate Change (IPCC Working Group III)

The IPCC AR5 Working Group III Summary for Policymakers (IPCC, 2014b) notes that total anthropogenic greenhouse gas emissions have continued to increase over 1970 to 2010 with larger absolute decadal increases toward the end of this period. Despite a growing number of climate change mitigation policies, annual emissions grew on average 2.2% per year from 2000 to 2010 compared with 1.3% per year from 1970 to 2000. Total anthropogenic greenhouse gas emissions were the highest in human history from 2000 to 2010. Globally, economic and population growth continue to be the most important drivers of increases in CO₂ emissions from fossil fuel combustion.

Limiting climate change will require substantial and sustained reductions of greenhouse gas emissions. The IPCC report considers multiple mitigation scenarios with a range of technological and behavioural options, with different characteristics and implications for sustainable development. These emissions scenarios are consistent with different levels of mitigation.

The IPCC report examines mitigation scenarios that would eventually stabilise greenhouse gases in the atmosphere at various concentration levels, and the expected corresponding changes in global temperatures. Mitigation scenarios where temperature change caused by anthropogenic greenhouse gas emissions can be kept to less than 2°C relative to pre-industrial levels involve stabilising atmospheric concentrations of carbon dioxide equivalent (CO₂-eq) at about 450 ppm in 2100. If concentration levels are not limited to 500 ppm CO₂-eq or less, temperature increases are unlikely to remain below 2°C relative to pre-industrial levels.

Without additional efforts to reduce emissions beyond those in place at present, scenarios project that global mean surface temperature increases in 2100 will be from 3.7 to 4.8°C compared to pre-industrial levels. This range is based on the median climate response, but when climate uncertainty is included the range becomes broader from 2.5 to 7.8°C.

In order to reach atmospheric greenhouse gas concentration levels of about 450 ppm CO₂-eq by 2100 (in order to have a likely chance to keep temperature change below 2°C relative to pre-industrial levels), anthropogenic greenhouse gas emissions would need to be cut by 40-70% globally by 2050 (compared with levels in 2010). Emissions levels would need to be near zero in 2100. The scenarios describe a wide range of changes to achieve this reduction in emissions, including large-scale changes in energy systems and land use.

Estimates of the cost of mitigation vary widely. Under scenarios in which all countries begin mitigation immediately, there is a single carbon price, and all key technologies are available, there will be losses of global consumption of 1-4% in 2030, 2-6% in 2050, and 3-11% in 2100.

Delaying mitigation efforts beyond those in place today through 2030 is estimated to substantially increase the difficulty in obtaining a longer term low level of greenhouse gas emissions, as well as narrowing the range of options available to maintain temperature change below 2°C relative to pre-industrial levels.

3 Background: New Zealand Climate Change – Science and Impacts

Published information about the expected impacts of climate change on New Zealand is summarised and assessed in the Australasia chapter of the IPCC Working Group II assessment report (Reisinger et al., 2014). Key findings from this chapter include:

The regional climate is changing. The Australasia region continues to demonstrate long-term trends toward higher surface air and sea surface temperatures, more hot extremes and fewer cold extremes, and changed rainfall patterns. Over the past 50 years, increasing greenhouse gas concentrations have contributed to rising average temperatures in New Zealand. Changing precipitation patterns have resulted in increases in rainfall for the south and west of the South Island and west of the North Island, and decreases in the northeast of the South Island and the east and north of the North Island. Some heavy rainfall events already carry the fingerprint of a changed climate, in that they have become more intense due to higher temperatures allowing the air to carry more moisture (Dean et al., 2013). Cold extremes have become rarer and hot extremes have become more common.

The region has exhibited warming to the present and is virtually certain to continue to do so. New Zealand mean annual temperature has increased by 0.09°C ($\pm 0.03^{\circ}\text{C}$) per decade since 1909.

Warming is projected to continue through the 21st century along with other changes in climate.

Warming is expected to be associated with rising snow lines, more frequent hot extremes, less frequent cold extremes, and increasing extreme rainfall related to flood risk in many locations. Annual average rainfall is expected to decrease in the northeast South Island and north and east of the North Island, and to increase in other parts of New Zealand. Fire weather is projected to increase in many parts of New Zealand. Regional sea level rise will very likely exceed the historical rate, consistent with global mean trends.

Uncertainty in projected rainfall changes remains large for many parts of New Zealand, which creates significant challenges for adaptation.

Impacts and vulnerability: Without adaptation, further climate-related changes are projected to have substantial impacts on water resources, coastal ecosystems, infrastructure, health, agriculture, and biodiversity. However, uncertainty in projected rainfall changes and other climate-related changes remains large for many parts of New Zealand, which creates significant challenges for adaptation.

Additional information about past New Zealand climate change can be found in (Mullan et al., 2016).

3.1 Sectoral Impacts

Some New Zealand sectors have the potential to benefit from projected changes in climate and increasing CO₂, including reduced winter mortality, reduced energy demand for winter heating, and forest and pasture growth in currently cooler regions.

Freshwater resources: In New Zealand, precipitation changes are projected to lead to increased runoff in the west and south of the South Island and reduced runoff in the northeast of the South Island, and the east and north of the North Island. Annual flows of eastward-flowing rivers with headwaters in the Southern Alps are projected to increase by 5-10% by 2040 in response to higher alpine precipitation. Most of the increases will occur in winter and spring, as more precipitation falls

as rain and snow melts earlier. Climate change will affect groundwater through changes in recharge rates and the relationship between surface waters and aquifers.

Natural ecosystems: Existing environmental stresses will interact with, and in many cases be exacerbated by, shifts in mean climatic conditions and associated changes in the frequency or intensity of extreme events, especially fire, drought, and floods. Ongoing impacts of invasive species and habitat loss will dominate climate change signals in the short to medium term. The rich biota of the alpine zone is at risk through increasing shrubby growth and loss of herbs, especially if combined with increased establishment of native species. Some cold water-adapted freshwater fish and invertebrates are vulnerable to warming and increased spring flooding may increase risks for braided river bird species.

Coastal and ocean ecosystems: The increasing density of coastal populations and stressors such as pollution and sedimentation from settlements and agriculture will intensify non-climate stressors in coastal areas. Coastal habitats provide many ecosystem services including coastal protection and carbon storage, which could become increasingly important for mitigation. Variability in ocean circulation and temperature plays an important role in local fish abundance, and this could change with climate-related oceanic changes. A strengthening East Auckland Current in northern New Zealand is expected to promote establishment of tropical or subtropical species that currently occur as vagrants, potentially changing the production and profit of both wild fisheries and aquaculture. Estuarine habitats will be affected by changing rainfall or sediment discharges, as well as connectivity to the ocean. Loss of coastal habitats and declines in iconic species will result in substantial impacts on coastal settlements and infrastructure from direct impacts such as storm surge, and will effect tourism. Changes in temperature and rainfall, and sea level rise, are expected to lead to secondary effects, including erosion, landslips, and flooding, affecting coastal habitats and their dependent species, for example loss of habitat for nesting birds.

Forestry: Warming is expected to increase *Pinus radiata* growth in the cooler south, whereas in the warmer north, temperature increases can reduce productivity, but CO₂ fertilisation may offset this. *Dothistroma* blight, a pine disease, has a temperature optimum that coincides with New Zealand's warmer, but not warmest, pine growing regions; under climate change, its severity is, therefore, expected to reduce in the warm central North Island but increase in the cooler South Island where it could offset temperature-driven improved plantation growth.

Agriculture: Projected changes in national pasture production for dairy, sheep, and beef pastures range from an average reduction of 4% across climate scenarios for the 2030s, to increases of up to 4% for two scenarios in the 2050s. Studies modelling seasonal changes in fodder supply show greater sensitivity in animal production to climate change and elevated CO₂ than models using annual average production, with some impacts expected even under modest warming. New Zealand agro-ecosystems are subject to erosion processes strongly driven by climate - greater certainty in projections of rainfall, particularly storm frequency, are needed to better understand climate change impacts on erosion and consequent changes in the ecosystem services provided by soils.

Energy supply, demand, and transmission: New Zealand's predominantly hydroelectric power generation is vulnerable to precipitation variability. Increasing winter precipitation and snow melt, and a shift from snowfall to rainfall will reduce this vulnerability as winter/spring inflows to main hydro lakes are projected to increase by 5-10% over the next few decades. Further reductions in seasonal snow and glacial melt as glaciers diminish, however, would compromise this benefit. Increasing wind power generation would benefit from projected increases in mean westerly winds

but face increased risk of damages and shutdown during extreme winds. Climate warming would reduce annual average peak electricity demands by 1-2% per degree Celsius across New Zealand.

Tourism: Changes in snow cover are likely to have a significant impact on the ski industry, but tourist numbers from Australia to New Zealand may increase due to the rapid reduction in snow cover in Australia, and the greater perceived scenic attractiveness of New Zealand. Warmer and drier conditions mostly benefit tourism but wetter conditions and extreme climate events undermine tourism.

4 Present Climate of Horizons Region

The climate of the Horizons Region is a reflection of the general disturbed westerly airflow with interspersed anticyclones, modified in specific places by the local topography (Chappell, 2012). Much of the region has relatively few climatic extremes, except in the higher elevation areas around the Central Plateau. The rainfall is usually adequate for pasture growth, except on occasions in the summer, and temperatures have a relatively small range. Summers are warm and frosts frequent in sheltered inland areas during winter. The weather is often cloudy about the hills, but sunshine hours increase toward the west coast. Except at higher elevations, snow and hail are rare occurrences, although fog occurs at times in coastal areas. The prevailing air flow is from the westerly quarter, and except during the passage of the occasional depression, or when a depression of tropical origin passes to the east of the North Island, the day-to-day weather conditions are not severe.

More detailed information about Horizon Region's present climate can be found in Chappell (2012).

4.1 Spatial Patterns in Horizons Region's Climate

The spatial variation in annual average temperature over the Horizons Region is shown in Figure 4-1. Figure 4-2 shows the spatial pattern of annual total rainfall, and also the median seasonal total rainfalls. Temperature varies with elevation, with the coolest mean annual temperatures of the Horizons Region experienced in the Central Plateau and the northern Ruahine Ranges. Mean annual temperatures are highest on the low elevation plains near the coast. Annual rainfall varies significantly throughout the Horizons Region. The high elevation areas of the Central Plateau, Ruahine Ranges, Tararua Ranges, as well as the hill country to the west of Taumarunui, receive the most rain in the region, due to orographic rainfall occurring from the prevailing westerly winds. These areas receive more than 2000 mm per year, on average. The driest part of the region is a triangle bounded by Whanganui, Levin, and Palmerston North, and also the area around Taihape. These parts receive around 850 mm of rainfall per year, on average.




Figure 4-1: Annual average temperature for the Horizons region (median for 1981-2010). ©NIWA.




Figure 4-2: Annual rainfall for the Horizons region (median for 1981-2010). ©NIWA.




Figure 4-2 continued: Seasonal rainfall totals (medians for spring, summer, autumn, and winter).
©NIWA.

4.2 Temporal Variability in Horizon Region's Climate

4.2.1 Temperature




Figure 4-3: Homogenised annual temperature time series for Whanganui from 1938 to 2014. The purple line removes the year-to-year variability and shows an upward long-term trend, and the grey shading shows the 95% confidence interval.

There is significant year-to-year variability in Horizon Region's climate. For example, Figure 4-3 shows the average annual temperature for Whanganui from 1938 to 2014. There are substantial differences between years, with some years having temperatures over 1.5°C different to others.

The temperature trend at Whanganui from 1938 to 2014 (shown on Figure 4-3), is 0.95 ± 0.41 °C. For comparison, the New Zealand national temperature¹ increased by 1.07 ± 0.40 °C over the same period. A likely explanation for the overall increase in average temperatures over this period is due to increasing concentrations of anthropogenic greenhouse gases, whereas the short-term variability is due to natural causes, such as the El Niño-Southern Oscillation and volcanic eruptions, together with random year-to-year fluctuations ("climate noise").

¹ <https://www.niwa.co.nz/our-science/climate/information-and-resources/nz-temp-record/seven-station-series-temperature-data>

4.2.2 Rainfall

As shown in Figure 4-4, there is also substantial variability in annual rainfall totals. At Whanganui, rainfall varies from around 600 mm per year to over 1200 mm, and Palmerston North rainfall varies from about 700 to 1100 mm, but there is no long term trend observed.




Figure 4-4: Annual rainfall total (mm) at Whanganui from 1890 to 2014 (top) and Palmerston North from 1944 to 2013 (bottom). Note change in site for Palmerston North. Metadata about climate stations can be found at <http://cliflo.niwa.co.nz>


4.2.3 Drought

Horizons Region has experienced numerous droughts, and due to the importance of primary production to the region, the occurrence of drought is of major concern. In this report, the measure of drought used is 'potential evapotranspiration deficit' (PED). Evapotranspiration is the combined loss of soil water by transpiration through plants and evaporative loss from the soil and other

surfaces. Days when water demand is not met, and pasture growth is reduced, are often referred to as days of potential evapotranspiration deficit. PED, in units of mm, can be thought of as the amount of rainfall needed in order to keep pastures growing at optimum levels.

The following plots (Figure 4-5 to Figure 4-8) show PED accumulations over growing years (July-June) through the historical record for the sites chosen. The higher the PED accumulation, the drier the soils were during that year. Note the change in scale between sites.

The highest PED accumulation for Taumarunui (Figure 4-5) occurred in 1969-70, with a secondary maximum in 2007-08. For most years, Taumarunui is not at risk of drought (i.e. high PED accumulations), because of its wetter climate compared with other parts of the region (Figure 4-2). However, the temporal variability in PED accumulation from year to year is quite variable, so at times, Taumarunui may be at risk of drought following a very wet year (i.e. no PED accumulation).



Copyright NIWA (C) 2016

Figure 4-5: PED accumulation from 1947-2015 (July-June years) for Taumarunui (agent number 2250).

Whanganui has experienced numerous years when high PED accumulations have been recorded (Figure 4-6). The highest accumulation occurred during 1976-77 and 2015-16, when over 500mm of PED accumulation was recorded, and 2007-08 recorded just under 500mm of accumulated PED. Compared with Taumarunui, Whanganui has a higher incidence of high PED accumulation, and therefore a higher risk of drought. Whanganui also has a much drier climate than Taumarunui (Figure 4-2).




Figure 4-6: PED accumulation from 1939-2015 (July-June years) for Wanganui, Spriggens Park EWS (agent number 3715).

Palmerston North (Figure 4-7) has a highly variable time series of PED accumulation, from almost no accumulation in some years to over 450mm accumulated in other years. The year with the highest PED accumulation was 1975-76.




Figure 4-7: PED accumulation from 1940-2015 (July-June years) for Palmerston N (agent number 3238) and Palmerston North EWS (agent number 21963) - combined series.

Like Palmerston North, Dannevirke has a highly variable time series of PED accumulation. The highest accumulations of PED occurred in 1977-78, 1986-87, and 1997-98.




Figure 4-8: PED accumulation from 1950-2015 (July-June years) for Dannevirke (agent number 2534).

4.2.4 Extreme Rainfall Events

High intensity rainfall is a concern in Horizons Region, due to flooding and soil erosion which impacts rivers, land-based primary industries, and urban settlements. Griffiths (2006) showed that during the periods 1930-2004 and 1950-2004, there were increases in the annual mean and annual extreme daily rainfall in the west of the North Island (west of a line from Wellington to Waipoua to Hamilton, i.e. including Horizons Region), and decreases to the east of that line. However, as shown in Figure 4-9, there is no observable trend in the number of days with >2mm, >5mm and >10mm of rain at Whanganui, Spriggens Park EWS climate station from 1938 to 2014. The three rainfall thresholds show a similar pattern.




Figure 4-9: Number of days per year with >2mm, >5mm and >10mm rain at Whanganui, Spriggens Park EWS site.

4.2.5 Growing Degree Days

The departure of mean daily temperature above a base temperature which has been found to be critical to the growth or development of a particular plant is a measure of the plant's development on that day. The sum of these departures then relates to the maturity or harvestable state of the crop. Thus, as the plant grows, updated estimates of harvest time can be made. These estimates have been found to be very valuable for a variety of crops with different base temperatures. Degree-day totals indicate the overall effects of temperature for a specified period, and can be applied to agricultural and horticultural production. Growing degree-days express the sum of daily temperatures above a selected base temperature that represent a threshold of plant growth. Figure 4-10 shows the median annual growing degree-day totals for base temperatures 5°C and 10°C for Horizons Region.




Figure 4-10: Median annual growing degree-days in Horizons Region, 1981-2010. Left: Base 5°C, Right: Base 10°C.

Air temperatures in Horizons Region have increased over the past century (Figure 4-3). As the calculation of growing degree-days is inherently dependent on temperature, one would expect to see an upward trend in the number of growing degree-days also.

4.3 Natural factors causing fluctuation in climate patterns over New Zealand

Much of the variation in New Zealand's climate is random and lasts for only a short period, but longer term, quasi-cyclic variations in climate can be attributed to different factors. Three large-scale oscillations that influence climate in New Zealand are the El Niño-Southern Oscillation, the Interdecadal Pacific Oscillation, and the Southern Annular Mode (Ministry for the Environment, 2008a).

4.3.1 The effect of El Niño and La Niña

The El Niño-Southern Oscillation (ENSO) is a natural mode of climate variability that has wide-ranging impacts around the Pacific basin (Ministry for the Environment, 2008a). The oscillation involves movement of warm ocean water from one side of the Pacific to the other, and the movement of rainfall across the Pacific associated with this warm water.

In an El Niño event, easterly trade winds weaken and warm water ‘spills’ across the Pacific towards the east, accompanied by higher rainfall than normal in the central-east Pacific. A La Niña event is essentially the opposite of this and is an intensification of ‘normal’ conditions, where the warm ocean waters remain over the western Pacific and the trade winds strengthen.

El Niño events occur on average 3 to 7 years apart, typically becoming established in April or May and persisting for about a year thereafter. The Southern Oscillation Index, or SOI, uses the pressure difference between Tahiti and Darwin to determine the state and intensity of ENSO. Persistence of about -1 signifies El Niño events, whereas +1 signifies La Niña (Figure 4-11).




Figure 4-11: Time series of the Southern Oscillation Index from 1930 to 2016. Blue shades are indicative of La Niña periods and red shades are indicative of El Niño periods.

The effects of El Niño and La Niña are most clearly observed in the tropics, but impacts are well-recognised in New Zealand also. In El Niño events, the weakened trade winds cause New Zealand to experience a stronger than normal south-westerly airflow. This generally brings lower seasonal temperatures to the country and drier than normal conditions to the north and east of New Zealand.

In La Niña conditions, the strengthened trade winds cause New Zealand to experience more north-easterly airflow than normal, higher temperatures, and wetter conditions in the north and east of the North Island. In the South Island higher pressures are often dominant, which can cause drought conditions there. Therefore, drought conditions can persist in either El Niño or La Niña phases in the South Island. Figure 4-12 shows average summer rainfall anomalies in New Zealand associated with El Niño and La Niña conditions. However, individual ENSO events may have significantly different rainfall patterns to those pictured.




Figure 4-12: Average summer percentage of normal rainfall during El Niño (left) and La Niña (right). ©NIWA

From Figure 4-12 it is evident that on average summer rainfall for the Manawatu Plains and the surrounding area is near normal during El Niño periods and slightly below normal during La Niña periods. The Ruapehu District and the eastern part of Horizons Region (east of the Ruahine and Tararua Ranges) receives below normal rainfall during El Niño periods.

According to the IPCC Assessment Report from Working Group I (IPCC, 2013), precipitation variability relating to ENSO will likely intensify due to increased moisture availability in the atmosphere. There is high confidence that ENSO will remain the dominant mode of natural climate variability in the 21st century. However, variations in the amplitude and spatial pattern of ENSO are large and therefore any specific projected changes in ENSO remain uncertain at this stage.

4.3.2 The effect of the Interdecadal Pacific Oscillation

The Interdecadal Pacific Oscillation, or IPO, is a large-scale, long period oscillation that influences climate variability over the Pacific Basin including New Zealand (Salinger et al., 2001). The IPO operates at a multi-decadal scale, with phases lasting around 20 to 30 years (Figure 4-13). During the positive phase of the IPO, sea surface temperatures around New Zealand tend to be lower, and westerly winds stronger, with the opposite occurring in the negative phase.




Figure 4-13: The Interdecadal Pacific Oscillation (IPO) index. Positive values indicate periods when stronger-than-normal westerlies occur over New Zealand, with more anticyclones than usual over northern New Zealand. Negative values indicate periods with more northeasterlies than normal over northern regions of the country. Vertical axis is the IPO index, and horizontal axis is the year.

New Zealand's climate appears to be affected by the long-term IPO cycle. The increase in New Zealand-wide temperatures around 1950 occurred shortly after the change from positive to negative phase of the IPO. In addition, the switch from negative to positive phase in 1977-78 coincided with significant rainfall changes (Ministry for the Environment, 2008a).




Figure 4-14: Percentage change in average annual rainfall, for the 1978-1998 period compared to the 1960-1977 period. (Note: From 1978-98 the IPO was in its positive phase, compared to the previous 18 years when the IPO was negative. Any local rainfall response due to global warming would also be contained within this pattern of rainfall trends). ©NIWA.

Figure 4-14 suggests that periods of positive IPO (which generally coincide with increased El Niño activity) tend to be drier, on average, for most of Horizons Region south of Ruapehu District. The IPO has been in a negative phase since 1999.

During the periods 1930-2004 and 1950-2004, a trend to increases in mean and extreme 1-day rainfall was generally observed in the west of both the North Island and South Island² (Griffiths, 2006). Griffiths suggests this results from a trend to increased westerly circulation across New Zealand between 1950 and 2004. This trend is consistent with enhanced warming since 1950 (as predicted by climate change modelling); the stronger IPO westerly phase since 1977; the increased frequency of El Niño events since 1977; or a mixture of all these considerations.

In Horizons Region, there is some evidence of changes in river flows in the Manawatu catchment with changes in the IPO, with positive phases of the IPO (i.e. 1978-1999) associated with lower flow conditions than negative phases (i.e. 1945-1977) (Henderson and Dietrich, 2007).

4.3.3 The effect of the Southern Annular Mode

The Southern Annular Mode (SAM) is a hemispheric atmospheric wave centred on the South Pole that affects New Zealand's climate in terms of westerly wind strength and storm occurrence (Kidston et al., 2009). In its positive phase, the SAM is associated with relatively light winds and more settled weather over New Zealand, with stronger westerly winds further south towards the pole. In contrast, the negative phase of the SAM is associated with unsettled weather over New Zealand and stronger westerly winds, whereas wind and storms decrease towards Antarctica.

In contrast to the longer-lived oscillations of ENSO and the IPO, each phase of the SAM may only last for a number of weeks before switching to the opposite phase. The phase and strength of the SAM is influenced by the size of the ozone hole, with the past increase in ozone depleting substances giving rise to a positive trend in the phase of the SAM (Thompson et al., 2011). However, with the recovery of the ozone hole and reduction of ozone-depleting substances projected into the future, the trend of summertime SAM phases is expected to become more negative and stabilise slightly above zero (i.e., it is expected that there will be slightly more positive SAM phases than negative phases. Note that the phases of the SAM are defined relative to the historical climate). However, increasing concentration of greenhouse gases in the atmosphere will have the opposite effect, of an increasing positive trend in summer and winter SAM phases, i.e. there will be more positive phases than negative phases into the future (Figure 4-15). The net result for SAM behaviour, as a consequence of both ozone recovery and greenhouse gas increases, is therefore likely to be relatively little change from present by 2100.

² The opposite behaviour – i.e. a trend to decreases in mean and extreme rainfall – was observed in the east of the North and South Islands.




Figure 4-15: Time series of the Southern Annular Mode from transient experiments forced with time-varying ozone-depleting substances and greenhouse gases. a. Forcing with ozone-depleting substances; b. forcing with greenhouse gases. The SAM index is defined as the leading principal component time series of 850-hpa Z anomalies 20-90°S; positive values of the index correspond to anomalously low Z over the polar cap, and vice versa. Lines denote the 50-year low-pass ensemble mean response for summer (DJF, solid black) and winter (JJA, dashed blue). Grey shading denotes +/- one standard deviation of the three ensemble members about the ensemble mean. The long-term means of the time series are arbitrary and are set to zero for the period 1970-1975. Past forcings are based on observational estimates; future forcings are based on predictions. After Thompson et al. (2011).

4.4 New Zealand Sea Level Trends and Variability

According to the IPCC AR5 Working Group I, global mean sea level rose by 0.19 ± 0.02 m from 1901 to 2010 (IPCC, 2013). Sea level rise around New Zealand is comparable to the global average, being approximately 0.17 ± 0.1 m for the 20th century (Reisinger et al., 2014).

Along with the long-term positive trend in sea level, there are short-term variations as well (Figure 4-16 and Table 4-1). Seasonal (annual), El Niño-Southern Oscillation (ENSO, 3-7 year), and Interdecadal Pacific Oscillation (IPO, 20-30 year) variations can cause fluctuations of up to about ± 0.25 m in background sea levels for short periods. For example during El Niño phases, sea levels around New Zealand tend to be depressed, and during La Niña phases sea levels around the country tend to be higher. The IPO in its negative phase tends to increase sea levels around the North Island by around 0.06 m above the background sea level rise.

Storm surge can also temporarily increase sea level over 1-3 days. Storm surge occurs due to a reduction in atmospheric pressure (inverse barometer effect) and the influence of the wind on the sea surface. In a New Zealand context, maximum storm surge on the open coast is unlikely to be more than 1 m, but can be higher in estuarine and harbour settings. Wave conditions also affect localised water levels where inshore of the wave breaker zone, water levels are set-up. This is a

localised phenomenon and can be highly variable along even a short stretch of coastline, being dependent on the wave conditions and configurations of offshore sandbars and beach slope.




Figure 4-16: Relative sea-level rise at four main New Zealand ports, 1900-2007. Modified after Hannah and Bell (2012). Annual MSL quality assurance undertaken by Prof. John Hannah and NIWA – data sourced originally from the port companies and obtained from the Land Information NZ archives. Each local vertical datum (LVD) was established from tide-gauge measurements in the earlier part of last century for years listed in Table 1 of Hannah & Bell (2012). Note: the plot shows the increase in annual MSL since the zero MSL datums were established from measurements in the earlier part of the 1900s, where the average passes through the zero line.

Table 4-1: Historical relative sea-level rise rates. Source: Hannah and Bell (2012). The SLR rates are relative to the local landmass at the sea-level gauge locations (and implicitly include vertical landmass movement).

Location	Historical rate of sea-level rise (mm yr^{-1})
Auckland	1.5 ± 0.1
Wellington	2.0 ± 0.2
Lyttelton	1.9 ± 0.1
Dunedin	1.3 ± 0.1

5 Projections of Horizons Region's Future Climate

Horizons Region's future climate will be influenced by a combination of the effects of anthropogenic climate change (increasing global concentrations of greenhouse gases, Section 2) plus the natural year-to-year and decade-to-decade variability resulting from "climate noise" and features such as the El Niño-Southern Oscillation (ENSO), the Interdecadal Pacific Oscillation (IPO), and the Southern Annular Mode (SAM), discussed in Section 4. This section first outlines the projected changes due to anthropogenic climate change in Horizons Region, and then returns to the issue of natural variability. Note that the projected changes use 20-year averages, which will not entirely remove effects of natural variability.

Predicting future changes in climate due to anthropogenic activity is made difficult because (a) predictions depend on future greenhouse gas concentrations, which in turn depend on global greenhouse gas emissions driven by factors such as economic activity, population changes, technological advances and policies for sustainable resource use, and (b) even for a specific future trajectory of global greenhouse gas emissions, different climate models predict somewhat different amounts of climate change.

This has been dealt with by the Intergovernmental Panel on Climate Change through consideration of 'scenarios' describing concentrations of greenhouse gases in the atmosphere associated with a range of possible economic, political, and social developments during the 21st century, and by considering results from several different climate models for a given scenario. In the 2013 IPCC Fifth Assessment Report, the atmospheric greenhouse gas concentration component of these scenarios are called Representative Concentrations Pathways (RCPs).

In Sections 5.1 and 5.2, global climate model output based on two RCPs has been downscaled to produce future projections for temperature and precipitation for the Horizons Region. The RCPs are based on 21st century climate policies, and thus differ from the previous IPCC SRES emissions scenarios and their 'no-climate policy' (IPCC, 2013). RCP 4.5 is a low-mid-range emissions scenario, which is also called a 'stabilisation' scenario where radiative forcing stabilises by 2100. RCP 8.5 is a scenario with very high greenhouse gas emissions, and radiative forcing continues to increase beyond 2100. Each RCP provides spatially-resolved data sets of land use change and sector-based emissions of air pollutants, and it specifies annual greenhouse gas concentrations and anthropogenic emissions up to 2500 (although this report only considers changes to 2100). RCPs are based on a combination of integrated assessment models, simple climate models, atmospheric chemistry and global carbon cycle models.

NIWA has used climate model data from the IPCC Fifth Assessment Report (IPCC, 2013) to update climate change scenarios for New Zealand, through both a regional climate model (dynamical) and statistical downscaling process. The dynamical and statistical downscaling processes are described in detail in an updated climate guidance manual prepared for the Ministry for the Environment (Mullan et al., 2016).

5.1 Horizons Climate Change Temperature Projections

The magnitude of the temperature change projections varies with the RCP and also with the climate models used. In this report, downscaling of two RCPs has been carried out to show the differences in temperature and precipitation projections for a stabilisation emissions scenario (RCP 4.5) and a high emissions scenario (RCP 8.5).

Figure 5-1 shows the seasonal patterns of projected temperature increase over the Horizons Region and surrounding areas for 2040 for the RCP 4.5 scenario, where the temperature changes of 37 climate models have been averaged together. Figure 5-2 shows corresponding patterns for 2090. Figure 5-3 shows the seasonal patterns of projected temperature increase for 2040 for the RCP 8.5 scenario, where the temperature changes of 41 climate models have been averaged together, and Figure 5-4 shows the corresponding patterns for 2090. These nominal years represent the mid-points of bi-decadal periods: 2040 is the average over 2031-2050, and 2090 the average over 2081-2100. All maps show changes relative to the baseline climate of 1986-2005.




Figure 5-1: Projected seasonal temperature changes at 2040 (2031-2050 average). Relative to 1986-2005 average, for the IPCC RCP 4.5 scenario, averaged over 37 climate models. ©NIWA.




Figure 5-2: Projected seasonal temperature changes at 2090 (2081-2100 average). Relative to 1986-2005 average, for the IPCC RCP 4.5 scenario, averaged over 37 climate models. ©NIWA.




Figure 5-3: Projected seasonal temperature changes at 2040 (2031-2050 average). Relative to 1986-2005 average, for the IPCC RCP 8.5 scenario, averaged over 41 climate models. ©NIWA.




Figure 5-4: Projected seasonal temperature changes at 2090 (2081-2100 average). Relative to 1986-2005 average, for the IPCC RCP 8.5 scenario, averaged over 41 climate models. ©NIWA.

Figures 5-1 and 5-2 show projected future warming in the Horizons Region of approximately 0.75°C – 1.0°C by 2040 for the RCP 4.5 scenario, when averaged over the 37 climate models analysed by NIWA. Figures 5-3 and 5-4 show projected future warming of approximately 1.0°C – 1.25°C by 2040 for the RCP 8.5 scenario, when averaged over 41 climate models. A slight acceleration in warming is projected for the second 50 years of the 21st century compared to the first 50 years, and inland areas tend to warm more (compared to the baseline) than coastal areas. Some models give less warming and others give a faster rate of warming (IPCC, 2013). The full range of model-projected warming is given in Table 5-1. The temperature ranges are relative to the baseline period 1986-2005 (as used by IPCC). Hence the projected changes at 2040 and 2090 should be thought of as 45-year and 95-year trends.

Table 5-1: Projected changes in seasonal and annual mean temperature (in °C) for the Horizons Region for 2040 and 2090. Changes are relative to the baseline period, 1986-2005. The changes are given for all four RCPs (2.6, 4.5, 6.0, 8.5), where the ensemble-average is taken over (23, 37, 18, 41) models, respectively. The first number is the ensemble average, with the bracketed numbers giving the range (5th and 95th percentile). After Mullan et al. (2016).

Period	RCP	Summer	Autumn	Winter	Spring	Annual
2040	RCP 8.5	1.1 (0.5, 1.8)	1.1 (0.7, 1.6)	1.1 (0.7, 1.5)	0.9 (0.4, 1.3)	1.1 (0.6, 1.6)
	RCP 6.0	0.9 (0.3, 1.6)	0.9 (0.3, 1.2)	0.8 (0.3, 1.2)	0.7 (0.2, 1.1)	0.8 (0.3, 1.1)
	RCP 4.5	0.9 (0.4, 1.5)	0.9 (0.4, 1.4)	0.9 (0.6, 1.2)	0.8 (0.3, 1.2)	0.9 (0.5, 1.2)
	RCP 2.6	0.7 (0.2, 1.3)	0.8 (0.3, 1.2)	0.7 (0.3, 1.1)	0.6 (0.3, 1.0)	0.7 (0.3, 1.1)
2090	RCP 8.5	3.3 (2.3, 4.7)	3.2 (2.3, 4.5)	3.2 (2.4, 4.1)	2.7 (1.9, 3.5)	3.1 (2.2, 4.4)
	RCP 6.0	1.9 (1.0, 3.6)	1.9 (1.0, 2.9)	1.9 (1.2, 2.8)	1.6 (1.0, 2.3)	1.8 (1.1, 2.9)
	RCP 4.5	1.5 (0.7, 2.7)	1.5 (0.8, 2.2)	1.5 (0.9, 2.1)	1.3 (0.7, 1.9)	1.4 (0.9, 2.1)
	RCP 2.6	0.7 (0.1, 1.4)	0.7 (0.2, 1.5)	0.7 (0.3, 1.3)	0.6 (0.2, 1.1)	0.7 (0.4, 1.3)

The seasonal and annual ensemble average projection (the number outside the brackets) in Table 5-1 is the temperature increase averaged over all 23 models for RCP 2.6, 37 models for RCP 4.5, 18 models for RCP 6.0, and 41 models for RCP 8.5 analysed by NIWA. The bracketed numbers give the range (5th and 95th percentile) for each RCP for each season and the annual projection.

By 2040 (2031-2050, relative to 1986-2005), annual average temperatures are projected to increase by between 0.7°C (RCP 2.6) and 1.1°C (RCP 8.5). Summer, autumn and winter have similar projections for warming, and the least warming is projected for spring. By 2090 (2081-2100, relative to 1986-2005), annual average temperatures are projected to increase by between 0.7°C for RCP 2.6 and 3.1°C for RCP 8.5. As for 2040, summer, autumn and winter have similar projections for warming, and the least warming is projected for spring. Note that the mitigation scenario (RCP 2.6) temperature change for 2090 is less than the change for 2040 in autumn and no change is observed for summer, winter, and spring, whereas all other emissions scenarios show increased warming at 2090 relative to 2040.

5.1.1 Projections of Growing Degree-Days

As discussed in Section 4.2.5, the calculation of growing degree-days is useful to primary industry in terms of monitoring plant growth and planning harvests. Due to projected mean temperature increases across the Horizons Region, as discussed in Section 5.1, it is expected that the annual total of growing degree-days will increase, with larger increases expected in the seasons that are likely to experience greater warming (i.e. summer, autumn, and winter). At present, future projections of Growing Degree Days based on the IPCC AR5 model data have not been performed for New Zealand.

5.1.2 Water Temperature Trends

Understanding potential future changes in river water temperature is important due to the impact of water temperature on freshwater ecosystems. Some invertebrate and fish species have narrow tolerance ranges with regard to temperature, so water temperature increases influenced by climate change-related air temperature increases could have a detrimental impact on these species. Preliminary modelling work has been carried out which suggests water temperature in New Zealand rivers will increase with projected climate change-induced air temperature increases (Doug Booker,

NIWA, pers. comm.). The amount of warming will vary depending on river elevation, catchment size, water source (e.g. snow melt or not) and so on.

According to the IPCC (Reisinger et al., 2014), there is high confidence that climate change is already affecting the oceans around New Zealand, with warming observed in the Tasman Sea off northern New Zealand. The western Tasman Sea is warming at a rate of four times the global average due to changes in the East Australian Current (Royal Society of New Zealand, 2016).

5.2 Projections for Frosts and Hot Days under Climate Change

As the seasonal mean temperature increases over time, we also expect to see changes in temperature extremes. In general, an increase in high temperature extremes, and a decrease in low temperature extremes is expected. Natural variability, of course, will continue to influence the climate of particular years, and the specific time variation of this variability cannot be predicted by the climate models due to the chaotic interactions that affect development of individual weather systems and larger-scale climate modes (such as El Niño events) (Mullan et al., 2016).

For this report, high temperature extremes (i.e. ‘hot days’) are considered as the number of days per year of 25°C or above, and low temperature extremes (i.e. ‘cold nights’ or frosts) are considered as the number of nights per year of 0°C or below. These extremes were determined by adding the monthly statistically downscaled temperature offsets to the daily VCSN³ maximum temperature (for ‘hot days’) and to daily VCSN minimum temperature (for ‘cold nights’) for each model, then counting the exceedances (greater than or equal to 25°C, or less than or equal to 0°C, for hot days and cold nights, respectively) for the selected RCP and time period. Finally the changes were averaged over the number of years (20) and the number of models (37 for RCP 4.5 and 41 for RCP 8.5).

Figure 5-5 shows the projected increase in the number of hot days per year at 2040 (2031-2050) and 2090 (2081-2100) relative to 1986-2005, for RCP 4.5 and RCP 8.5. At 2040 there is projected to be only a small increase in the number of hot days in higher elevations around the Central Plateau, with the coastal area south of Whanganui and the hill country around Taumarunui receiving the largest projected increase (an increase of 15-20 days under RCP 8.5 compared to 1986-2005). By 2090, the projected change in hot days is much larger, with an expected increase of 50-60 days under RCP 8.5 compared to 1986-2005 on the coast near and south of Whanganui and the hill country north towards Taumarunui.

³ Virtual Climate Station Network, a set of New Zealand climate data based on a 5 km by 5 km grid across the country. Data has been interpolated from ‘real’ climate station records (TAIT, A., HENDERSON, R., TURNER, R. & ZHENG, X. G. 2006. Thin plate smoothing spline interpolation of daily rainfall for New Zealand using a climatological rainfall surface. *International Journal of Climatology*, 26, 2097-2115.)




Figure 5-5: Projected increase in number of hot days per year ($T_{max} > 25^{\circ}\text{C}$) at 2040 & 2090 for RCP 4.5 (left panels) and RCP 8.5 (right panels), for Horizons Region. Projected change in hot days is relative to 1986-2005. The numbers on the scale refer to the *increase* in the number of hot days, e.g. the coast adjacent to Whanganui is projected to experience an increase in the number of hot days by 50-60 days by 2090 under RCP 8.5 (lower right panel). © NIWA.

Figure 5-6 shows the projected decrease in the number of cold nights (or frosts) per year by 2040 (2031-2050) and 2090 (2081-2100) for RCP 4.5 and RCP 8.5. The projected decrease in the number of cold nights at both 2040 and 2090 for both RCPs is most significant in the Central Plateau and higher elevation areas of the Ruahine and Tararua Ranges. By 2090 under RCP 8.5, the number of cold nights in the Central Plateau is expected to decrease by more than 50 nights per year compared to 1986-2005 (with a decrease of more than 80 nights per year projected for the alpine areas of the Tongariro National Park). By 2090 under RCP 8.5 along the coast and inland between Whanganui and Levin, the number of cold nights is projected to decrease by 5-15 nights per year compared to 1986-2005.




Figure 5-6: Projected decrease in number of cold nights per year ($T_{min} < 0^{\circ}\text{C}$) at 2040 and 2090 for RCP 4.5 (left) and RCP 8.5 (right), for Horizons Region. Projected change in cold nights is relative to 1986-2005. The numbers on the scale refer to the *decrease* in the number of cold nights, e.g. the area around Taihape is projected to experience a decrease in the number of cold nights by 40-50 days by 2090 under RCP 8.5 (lower right panel). © NIWA.

As shown in Table 5-2, projections suggest a substantial decrease in the annual number of cold nights (or frosts) on average across Horizons Region, with perhaps 28 fewer cold nights by 2090 under RCP 8.5 (a decrease from 36 days per year during 1986-2005 to 2 days per year during 2081-2100 under RCP 8.5). The number of hot days is projected to increase significantly, with perhaps 47 more hot days by 2090 under RCP 8.5 (an increase from 19 days per year during 1986-2005 to 66 days per year during 2081-2100 under RCP 8.5). The values in Table 5-2 are a calculated average over the Horizons Region for all VCSN points below 500 m altitude; hence the projected changes are lower (or higher) than some of the local-scale changes in Figure 5-5 and Figure 5-6.

Table 5-2: Projected changes in the number of hot days and cold nights (frosts) at 2040 and 2090 for Horizons Region. Model results from RCP 4.5 and RCP 8.5 are compared to the historical period (1986-2005). The projected changes were averaged over all VCSN points across the Horizons Region below 500 m elevation. After Mullan et al. (2016).

		Historical period days	2040 number of days	2040 change	2090 number of days	2090 change
Hot days	RCP 4.5	19	29	+10	37	+18
	RCP 8.5		31	+12	66	+47
Cold nights	RCP 4.5	18	11	-7	7	-11
	RCP 8.5		9	-11	2	-16

5.3 Horizons Climate Change Precipitation Projections

Precipitation (rain + snow) projections show more spatial variation than the temperature projections. Again, the magnitude of the projected change will scale up or down with the different RCPs, and will also differ between climate models. Figure 5-7 and Figure 5-8 show the projected seasonal patterns of precipitation change over the Horizons Region and surrounding areas at 2040 and 2090 for RCP 4.5 (averaging 37 climate models), and Figure 5-9 and Figure 5-10 show the same for RCP 8.5 (averaging 41 climate models).




Figure 5-7: Projected seasonal precipitation changes (in %) at 2040 (2031-2050 average). Relative to 1986-2005 average, for the IPCC RCP 4.5 scenario, averaged over 37 climate models. ©NIWA.




Figure 5-8: Projected seasonal precipitation changes (in %) at 2090 (2081-2100 average). Relative to 1986-2005 average, for the RCP 4.5 scenario, averaged over 37 climate models. ©NIWA.




Figure 5-9: Projected seasonal precipitation changes (in %) at 2040 (2031-2050 average). Relative to 1986-2005 average, for the IPCC RCP 8.5 scenario, averaged over 41 climate models. ©NIWA.




Figure 5-10: Projected seasonal precipitation changes (in %) at 2090 (2081-2100 average). Relative to 1986-2005 average, for the IPCC RCP 8.5 scenario, averaged over 41 climate models. ©NIWA.

The RCP 4.5 and RCP 8.5 projections to 2040 (Figure 5-7 and Figure 5-9) indicate slightly more rainfall in winter (up to 10% more) for the western part of Horizons Region. For the other seasons, the projected precipitation change is $\pm 5\%$. By 2090 for RCP 4.5 (Figure 5-8), more rainfall is projected for the hill country west of Raetihi (up to 15% more) in winter, but for the other seasons the projected precipitation change remains $\pm 5\%$.

For RCP 8.5 at 2090 in winter (Figure 5-10), the northern part of the region is projected to experience up to 20% more precipitation, whereas precipitation for the eastern part of the region is projected to decrease by up to 20% in some areas. In summer, the eastern part of Horizons Region is projected to be wetter compared with the baseline, with up to 15% more precipitation projected for some parts, whereas the western part of the region may experience a small decrease in summer precipitation.

The full range of model-projected precipitation change (in %) is given for a number of locations in Horizons Region (grid points chosen by Horizons Regional Council). Model-projected changes are given for 2031-2050 (2040) in Table 5-3 and for 2081-2100 (2090) in Table 5-4. The precipitation changes are relative to the baseline period 1986-2005. Hence the projected changes at 2040 and 2090 should be thought of as 45-year and 95-year trends.

Table 5-3: Projected changes in seasonal and annual precipitation (in %) between 1986-2005 and 2031-2050, for selected locations within the Horizons Regional Council area, as derived from statistical downscaling. The changes are given for four RCPs (8.5, 6.0, 4.5 and 2.6), where the ensemble-average is taken over (41, 18, 37, 23) models respectively. The values in each column represent the ensemble average, and in brackets the range (5th percentile to 95th percentile) over all models within that ensemble. © NIWA.

Region	Summer	Autumn	Winter	Spring	Annual
Taumarunui					
rcp 8.5	0 (-7, 11)	-1 (-11, 9)	8 (-8, 24)	1 (-9, 17)	2 (-3, 10)
rcp 6.0	-1 (-13, 9)	0 (-16, 12)	7 (-7, 21)	0 (-10, 8)	2 (-5, 10)
rcp 4.5	0 (-9, 9)	0 (-13, 11)	7 (-7, 19)	2 (-12, 12)	2 (-4, 8)
rcp 2.6	0 (-6, 11)	0 (-8, 11)	4 (-13, 18)	1 (-9, 11)	2 (-4, 9)
Pipiriki					
rcp 8.5	0 (-10, 10)	-1 (-11, 7)	8 (-8, 25)	2 (-9, 19)	2 (-3, 11)
rcp 6.0	-2 (-12, 11)	-1 (-14, 8)	8 (-7, 22)	0 (-10, 10)	2 (-5, 11)
rcp 4.5	0 (-11, 8)	-1 (-12, 8)	7 (-6, 19)	2 (-12, 13)	2 (-4, 8)
rcp 2.6	0 (-6, 10)	0 (-8, 8)	4 (-14, 18)	2 (-9, 13)	2 (-4, 9)
Whanganui					
rcp 8.5	0 (-10, 11)	-1 (-10, 7)	5 (-6, 16)	0 (-9, 14)	1 (-5, 7)
rcp 6.0	-1 (-11, 9)	0 (-11, 7)	6 (-5, 16)	0 (-7, 9)	1 (-4, 8)
rcp 4.5	0 (-9, 9)	-1 (-9, 8)	5 (-6, 15)	1 (-8, 10)	1 (-4, 6)
rcp 2.6	0 (-8, 11)	0 (-8, 5)	3 (-9, 16)	1 (-8, 9)	1 (-4, 7)
Bulls					
rcp 8.5	0 (-10, 11)	-1 (-11, 9)	6 (-6, 17)	1 (-9, 15)	2 (-5, 9)
rcp 6.0	-1 (-11, 10)	0 (-14, 10)	6 (-5, 18)	0 (-8, 10)	1 (-4, 9)
rcp 4.5	0 (-9, 10)	0 (-12, 10)	5 (-5, 15)	2 (-9, 11)	2 (-4, 6)
rcp 2.6	0 (-8, 11)	1 (-9, 9)	3 (-10, 16)	1 (-8, 10)	1 (-3, 7)
Foxton					
rcp 8.5	0 (-10, 11)	0 (-9, 9)	6 (-5, 19)	0 (-10, 13)	2 (-5, 9)
rcp 6.0	-1 (-11, 11)	1 (-12, 11)	6 (-5, 20)	0 (-9, 11)	1 (-5, 9)
rcp 4.5	0 (-10, 10)	1 (-10, 11)	6 (-5, 15)	1 (-8, 10)	2 (-4, 7)
rcp 2.6	0 (-8, 10)	1 (-8, 9)	4 (-10, 16)	1 (-8, 8)	1 (-3, 7)
Levin					
rcp 8.5	0 (-10, 10)	0 (-9, 9)	7 (-5, 20)	1 (-10, 14)	2 (-4, 9)
rcp 6.0	-1 (-10, 11)	1 (-11, 10)	6 (-5, 20)	0 (-8, 11)	2 (-5, 9)
rcp 4.5	0 (-10, 9)	1 (-9, 10)	6 (-5, 16)	1 (-8, 11)	2 (-4, 7)
rcp 2.6	0 (-7, 10)	1 (-8, 9)	4 (-10, 16)	1 (-7, 9)	1 (-3, 7)
Shannon					
rcp 8.5	0 (-11, 11)	0 (-10, 9)	6 (-5, 19)	1 (-9, 14)	2 (-5, 9)
rcp 6.0	-2 (-11, 12)	1 (-12, 11)	6 (-5, 19)	0 (-8, 11)	2 (-5, 10)
rcp 4.5	0 (-11, 10)	1 (-10, 11)	6 (-5, 16)	2 (-8, 11)	2 (-4, 7)
rcp 2.6	-1 (-8, 10)	1 (-8, 10)	4 (-10, 16)	1 (-7, 9)	1 (-3, 8)
Palmerston North					
rcp 8.5	0 (-9, 11)	0 (-10, 9)	6 (-6, 19)	1 (-8, 15)	2 (-4, 9)
rcp 6.0	-1 (-11, 9)	0 (-11, 10)	7 (-5, 19)	0 (-7, 10)	2 (-4, 10)
rcp 4.5	0 (-9, 11)	0 (-9, 10)	6 (-5, 16)	2 (8, 11)	2 (-3, 7)
rcp 2.6	0 (-8, 10)	1 (-8, 7)	4 (-11, 17)	1 (7, 10)	2 (-3, 8)
Taihape					
rcp 8.5	0 (-10, 9)	-1 (-9, 4)	5 (-6, 17)	1 (-8, 14)	1 (-3, 6)
rcp 6.0	1 (-8, 16)	0 (-9, 6)	6 (-6, 18)	1 (-7, 11)	2 (-3, 6)
rcp 4.5	0 (-8, 9)	-1 (-8, 6)	5 (-7, 16)	2 (-7, 10)	2 (-2, 7)
rcp 2.6	2 (-5, 12)	-1 (-8, 5)	3 (-10, 18)	1 (-8, 10)	1 (-4, 7)
Ohakune					
rcp 8.5	0 (-8, 11)	-1 (-11, 7)	8 (-8, 26)	2 (-10, 19)	2 (-3, 11)
rcp 6.0	-1 (-12, 9)	-1 (-14, 7)	8 (-8, 23)	1 (-8, 9)	2 (-4, 11)
rcp 4.5	0 (-10, 7)	-1 (-12, 7)	8 (-6, 20)	2 (-11, 14)	3 (-3, 7)

Region	Summer	Autumn	Winter	Spring	Annual
rcp 2.6	0 (-7, 10)	0 (-8, 7)	5 (-15, 19)	2 (-9, 14)	2 (-4, 10)
Waimiha	rcp 8.5	0 (-7, 11)	0 (-11, 9)	8 (-8, 23)	1 (-9, 16)
	rcp 6.0	-1 (-14, 8)	0 (-16, 14)	7 (-6, 20)	0 (-10, 7)
	rcp 4.5	0 (-9, 9)	0 (-13, 11)	7 (-7, 19)	1 (-12, 11)
	rcp 2.6	0 (-6, 10)	1 (-8, 12)	4 (-13, 18)	1 (-8, 10)
Ruahine Ranges	rcp 8.5	0 (-9, 8)	0 (-6, 6)	2 (-6, 9)	-1 (-8, 7)
	rcp 6.0	1 (-8, 13)	1 (-10, 6)	2 (-9, 11)	0 (-8, 7)
	rcp 4.5	0 (-8, 10)	0 (-8, 8)	2 (-9, 10)	1 (-7, 9)
	rcp 2.6	1 (-6, 10)	0 (-7, 5)	1 (-4, 9)	0 (-9, 9)
Tararua Ranges	rcp 8.5	0 (-11, 12)	1 (-9, 9)	6 (-5, 18)	1 (-12, 15)
	rcp 6.0	-2 (-11, 13)	1 (-11, 11)	6 (-7, 19)	0 (-10, 12)
	rcp 4.5	0 (-10, 9)	1 (-9, 11)	6 (-6, 16)	2 (-10, 12)
	rcp 2.6	-1 (-8, 10)	2 (-8, 11)	3 (-8, 14)	1 (-8, 11)
Eketahuna	rcp 8.5	0 (-16, 15)	-1 (-11, 9)	5 (-4, 16)	2 (-8, 16)
	rcp 6.0	-3 (-17, 17)	0 (-12, 7)	5 (-6, 17)	1 (-8, 11)
	rcp 4.5	-1 (-16, 12)	0 (-10, 9)	5 (-6, 15)	3 (-9, 13)
	rcp 2.6	-1 (-13, 11)	0 (-8, 7)	3 (-7, 14)	2 (-8, 12)
Dannevirke	rcp 8.5	0 (-11, 7)	0 (-8, 6)	0 (-7, 8)	0 (-7, 10)
	rcp 6.0	2 (-10, 15)	1 (-12, 8)	2 (-5, 8)	0 (-7, 9)
	rcp 4.5	0 (-8, 13)	0 (-7, 9)	1 (-7, 9)	2 (-5, 10)
	rcp 2.6	2 (-5, 10)	0 (-7, 5)	1 (-4, 9)	0 (-8, 10)
Pahiatua	rcp 8.5	0 (-9, 10)	-1 (-10, 8)	5 (-5, 15)	1 (-8, 15)
	rcp 6.0	-1 (-13, 9)	0 (-10, 8)	6 (-4, 16)	1 (-7, 10)
	rcp 4.5	0 (-11, 12)	0 (-8, 8)	5 (-7, 16)	2 (-8, 11)
	rcp 2.6	0 (-10, 10)	0 (-7, 6)	3 (-9, 16)	2 (-7, 10)
Akitio	rcp 8.5	0 (-17, 10)	2 (-10, 13)	-5 (-19, 4)	-3 (-14, 8)
	rcp 6.0	5 (-9, 25)	3 (-15, 13)	-3 (-16, 6)	0 (-10, 12)
	rcp 4.5	1 (-10, 16)	2 (-10, 14)	-5 (-16, 11)	0 (-11, 16)
	rcp 2.6	3 (-8, 14)	1 (-9, 11)	-2 (-14, 12)	-2 (-14, 10)
Owahanga	rcp 8.5	0 (-16, 10)	2 (-10, 13)	-5 (-18, 5)	-3 (-14, 8)
	rcp 6.0	4 (-9, 23)	3 (-15, 12)	-3 (-14, 6)	0 (-10, 11)
	rcp 4.5	1 (-10, 16)	2 (-10, 14)	-4 (-14, 10)	0 (-11, 14)
	rcp 2.6	3 (-8, 13)	1 (-9, 11)	-2 (-14, 12)	-2 (-13, 9)

Table 5-4: Projected changes in seasonal and annual precipitation (in %) between 1986-2005 and 2081-2100, for selected locations within the Horizons Regional Council area, as derived from statistical downscaling. The changes are given for four RCPs (8.5, 6.0, 4.5 and 2.6), where the ensemble-average is taken over (41, 18, 37, 23) models respectively. The values in each column represent the ensemble average, and in brackets the range (5 percentile to 95 percentile) over all models within that ensemble. © NIWA.

Region	Summer	Autumn	Winter	Spring	Annual
Taumarunui	rcp 8.5	2 (-15, 22)	-5 (-24, 10)	16 (-10, 36)	0 (-22, 14)
	rcp 6.0	2 (-22, 21)	2 (-22, 45)	15 (-6, 65)	5 (-16, 42)
	rcp 4.5	2 (-10, 13)	0 (-11, 12)	9 (-14, 28)	1 (-9, 12)
	rcp 2.6				3 (-7, 11)

Region		Summer	Autumn	Winter	Spring	Annual
	rcp 2.6	2 (-10, 15)	2 (-10, 12)	7 (-5, 24)	4 (-6, 18)	4 (-4, 15)
Pipiriki	rcp 8.5	0 (-26, 19)	-6 (-22, 7)	17 (-11, 37)	2 (-22, 17)	4 (-14, 14)
	rcp 6.0	0 (-29, 15)	-2 (-20, 13)	14 (-5, 39)	4 (-17, 18)	4 (-14, 14)
	rcp 4.5	2 (-12, 15)	-1 (-8, 9)	10 (-14, 29)	2 (-8, 14)	4 (-7, 11)
	rcp 2.6	3 (-8, 17)	2 (-9, 9)	7 (-6, 25)	5 (-6, 21)	4 (-3, 15)
Whanganui	rcp 8.5	0 (-20, 20)	-5 (-16, 7)	10 (-12, 28)	-1 (-19, 10)	1 (-11, 10)
	rcp 6.0	2 (-25, 35)	0 (-15, 23)	11 (-8, 43)	33 (-12, 18)	4 (-12, 30)
	rcp 4.5	2 (-13, 14)	0 (-7, 8)	6 (-10, 21)	1 (-8, 11)	2 (-5, 11)
	rcp 2.6	3 (-8, 14)	2 (-8, 6)	6 (-5, 20)	3 (-8, 14)	3 (-3, 13)
Bulls	rcp 8.5	-1 (-22, 19)	-4 (-19, 8)	12 (-11, 29)	1 (-18, 12)	2 (-12, 11)
	rcp 6.0	0 (-26, 13)	-1 (-19, 16)	10 (-7, 30)	3 (-13, 15)	3 (-13, 13)
	rcp 4.5	2 (-13, 15)	0 (-7, 10)	7 (-12, 22)	2 (-7, 12)	3 (-6, 10)
	rcp 2.6	3 (-9, 15)	2 (-8, 11)	6 (-5, 21)	4 (-7, 16)	4 (-3, 14)
Foxton	rcp 8.5	-1 (-23, 18)	-2 (-16, 10)	13 (-7, 32)	0 (-17, 11)	3 (-10, 11)
	rcp 6.0	0 (-26, 13)	1 (-16, 17)	11 (-5, 30)	2 (-12, 11)	3 (-12, 13)
	rcp 4.5	2 (-12, 15)	1 (-6, 9)	8 (-10, 23)	1 (-6, 10)	3 (-6, 10)
	rcp 2.6	3 (-9, 16)	3 (-7, 12)	6 (-5, 21)	3 (-8, 12)	3 (-3, 14)
Levin	rcp 8.5	0 (-23, 18)	-2 (-16, 10)	14 (-7, 34)	1 (-16, 13)	3 (-10, 12)
	rcp 6.0	0 (-26, 13)	1 (-15, 16)	11 (-5, 31)	2 (-11, 12)	4 (-12, 14)
	rcp 4.5	2 (-11, 16)	1 (-5, 9)	9 (-10, 23)	1 (-6, 11)	3 (-5, 11)
	rcp 2.6	3 (-9, 16)	3 (-6, 11)	6 (-5, 21)	3 (-7, 13)	4 (-3, 14)
Shannon	rcp 8.5	-1 (-27, 18)	-2 (-16, 10)	13 (-8, 32)	1 (-17, 13)	3 (-11, 12)
	rcp 6.0	-1 (-29, 12)	1 (-16, 17)	11 (-5, 30)	3 (-11, 14)	4 (-13, 14)
	rcp 4.5	2 (-13, 16)	1 (-6, 9)	8 (-10, 22)	2 (-6, 12)	3 (-6, 11)
	rcp 2.6	3 (-10, 17)	3 (-7, 12)	6 (-5, 21)	3 (-7, 14)	4 (-3, 14)
Palmerston North	rcp 8.5	0 (-17, 20)	-3 (-16, 8)	13 (-11, 31)	1 (-18, 13)	3 (-10, 11)
	rcp 6.0	0 (-22, 12)	0 (-15, 14)	11 (-6, 33)	3 (-12, 15)	4 (-11, 13)
	rcp 4.5	2 (-11, 13)	1 (-6, 8)	8 (-12, 23)	2 (-7, 12)	3 (-6, 11)
	rcp 2.6	2 (-8, 14)	2 (-7, 9)	6 (-6, 22)	4 (-7, 16)	4 (-3, 14)
Taihape	rcp 8.5	7 (-5, 28)	-5 (-15, 8)	11 (-14, 32)	-1 (-18, 12)	3 (-3, 11)
	rcp 6.0	4 (-6, 18)	-2 (-12, 9)	10 (-8, 30)	2 (-10, 17)	4 (-4, 12)
	rcp 4.5	2 (-5, 10)	-1 (-8, 6)	7 (-11, 23)	1 (-7, 13)	2 (-2, 10)
	rcp 2.6	0 (-11, 9)	1 (-8, 8)	6 (-5, 21)	4 (-4, 13)	3 (-4, 11)
Ohakune	rcp 8.5	1 (-19, 20)	-7 (-22, 7)	18 (-11, 38)	3 (-22, 19)	4 (-13, 14)
	rcp 6.0	0 (-25, 15)	-2 (-19, 11)	14 (-5, 41)	4 (-16, 19)	5 (-12, 14)
	rcp 4.5	2 (-11, 13)	-1 (-8, 8)	10 (-15, 29)	2 (-8, 15)	4 (-6, 12)
	rcp 2.6	2 (-8, 16)	1 (-10, 8)	7 (-6, 26)	5 (-7, 23)	4 (-3, 15)
Waimiha	rcp 8.5	2 (-18, 21)	-4 (-24, 11)	16 (-8, 36)	0 (-22, 13)	4 (-15, 14)
	rcp 6.0	1 (-24, 17)	0 (-22, 18)	13 (-6, 35)	2 (-16, 15)	4 (-13, 14)
	rcp 4.5	2 (-11, 13)	1 (-11, 11)	9 (-14, 27)	1 (-9, 12)	3 (-7, 10)
	rcp 2.6	2 (-9, 15)	2 (-10, 13)	7 (-5, 23)	4 (-6, 17)	4 (-4, 15)
Ruahine Ranges	rcp 8.5	4 (-9, 25)	-1 (-11, 8)	2 (-13, 13)	-4 (-17, 7)	0 (-6, 8)
	rcp 6.0	3 (-7, 16)	1 (-8, 8)	3 (-10, 19)	0 (-9, 10)	2 (-6, 10)

Region	Summer	Autumn	Winter	Spring	Annual
rcp 4.5	2 (-7, 10)	1 (-9, 7)	2 (-9, 11)	0 (-10, 9)	1 (-4, 6)
rcp 2.6	1 (-8, 9)	1 (-5, 8)	2 (-4, 9)	2 (-7, 10)	2 (-5, 9)
Tararua Ranges					
rcp 8.5	-1 (-31, 22)	-1 (-15, 11)	14 (-4, 29)	2 (-15, 16)	4 (-11, 12)
rcp 6.0	-1 (-31, 13)	1 (-15, 18)	10 (-5, 27)	3 (-13, 15)	4 (-13, 15)
rcp 4.5	2 (-12, 17)	2 (-6, 10)	8 (-7, 21)	2 (-6, 15)	4 (-5, 11)
rcp 2.6	3 (-10, 20)	3 (-8, 13)	5 (-3, 19)	3 (-6, 14)	4 (-3, 14)
Eketahuna					
rcp 8.5	-6 (-42, 20)	-5 (-17, 8)	12 (-4, 27)	3 (-17, 17)	2 (-13, 11)
rcp 6.0	-3 (-40, 17)	-1 (-16, 12)	9 (-6, 26)	4 (-11, 19)	3 (-15, 14)
rcp 4.5	1 (-19, 18)	0 (-7, 7)	7 (-7, 19)	3 (-6, 16)	3 (-6, 11)
rcp 2.6	5 (-12, 21)	2 (-8, 7)	5 (-3, 18)	5 (-6, 20)	4 (-3, 14)
Dannevirke					
rcp 8.5	5 (-8, 26)	-1 (-11, 9)	-1 (-14, 9)	-3 (-18, 12)	0 (-6, 7)
rcp 6.0	4 (-6, 15)	0 (-7, 9)	2 (-12, 17)	1 (-8, 13)	2 (-3, 10)
rcp 4.5	2 (-9, 13)	0 (-8, 7)	1 (-8, 10)	0 (-8, 11)	1 (-4, 5)
rcp 2.6	0 (-9, 11)	1 (-5, 10)	2 (-5, 9)	3 (-5, 12)	2 (-5, 9)
Pahiatua					
rcp 8.5	-1 (-16, 21)	-4 (-16, 8)	10 (-12, 27)	2 (-17, 13)	2 (-10, 10)
rcp 6.0	0 (-21, 11)	-1 (-14, 12)	9 (-8, 27)	3 (-10, 17)	3 (-10, 13)
rcp 4.5	2 (-11, 13)	0 (-7, 8)	7 (-10, 23)	2 (-7, 13)	3 (-5, 12)
rcp 2.6	2 (-7, 12)	2 (-7, 8)	6 (-4, 19)	4 (-6, 16)	4 (-3, 13)
Akitio					
rcp 8.5	12 (-7, 33)	7 (-8, 19)	-15 (-35, 2)	-9 (-27, 4)	-2 (-11, 11)
rcp 6.0	8 (-9, 20)	4 (-11, 15)	-8 (-32, 8)	-4 (-18, 11)	0 (-9, 9)
rcp 4.5	4 (-11, 15)	2 (-12, 13)	-6 (-26, 9)	-2 (-17, 10)	-1 (-13, 7)
rcp 2.6	-2 (-18, 9)	1 (-8, 12)	-2 (-10, 9)	0 (-11, 11)	-1 (-8, 5)
Owahanga					
rcp 8.5	10 (-8, 32)	7 (-7, 19)	-14 (-34, 2)	-9 (-26, 4)	-3 (-11, 11)
rcp 6.0	7 (-9, 19)	4 (-11, 15)	-7 (-30, 8)	-3 (-18, 10)	0 (-8, 9)
rcp 4.5	4 (-11, 16)	2 (-12, 13)	-5 (-25, 9)	-2 (-17, 9)	-1 (-12, 6)
rcp 2.6	-2 (-16, 9)	1 (-9, 12)	-1 (-10, 8)	0 (-10, 10)	-1 (-8, 5)

The seasonal and annual ensemble average projection (the number outside the brackets) in Table 5-3 and Table 5-4 is the precipitation increase (in %) for the given locations for 2040 and 2090, respectively, averaged over all 23 models for RCP 2.6, 37 models for RCP 4.5, 18 models for RCP 6.0, and 41 models for RCP 8.5 analysed by NIWA. The bracketed numbers give the range (5th and 95th percentile) for each RCP for each season and the annual projection.

For many locations, there is no clear seasonal and annual precipitation signal, even at 2090 under RCP 8.5. The ensemble-average is often less than ±5%, with the model range (the 5th and 95th percentile values) varying between quite large (>10%) increases and decreases. By 2040 (2031-2050, relative to 1986-2005), winter is the season with the most precipitation change, with a small increase in the ensemble-average at most locations (up to 8% across the different locations and RCPs). Akitio and Owahanga are the only locations to show a negative precipitation change by 2040.

By 2090 (2081-2100, relative to 1986-2005), there is a clearer precipitation signal at most locations. For most locations, winter is still the season with the most precipitation change, with some locations projecting increases in the ensemble-average at around 15% (Ohakune has the highest projected winter precipitation increase of 18% at RCP 8.5). However, the direction of projected precipitation

change is not uniform across the region; precipitation for Akitio and Owahanga is projected to decrease by ~15% by 2090 under RCP 8.5.

5.4 Temperature and precipitation projection comparisons within RCPs

The average picture of projected temperature and rainfall changes in the tables and maps in Sections 5.1 to 5.3 obscures significant variations between the individual models run under each RCP on the projected seasonal changes. Figure 5-11 and Figure 5-12 show seasonal temperature projections from all the models individually averaged over the Horizons Region for 2040 and 2090, respectively. The coloured vertical bars, and inset stars, show the individual models, so the complete range is displayed (unlike Tables 5-1, 5-3, and 5-4 where the 5th to 95th percentile range has been calculated). Figure 5-11 and Figure 5-12 show an excellent way of not only demonstrating the difference with season and RCP, but also the range of model sensitivity. The black stars within each vertical bar represent the results of the 6 Regional Climate Model (RCM) simulations; the RCM projections tend to be in the lower half of the statistically-downscaled results, owing to the bias-correction applied to the raw RCM output.

For 2040 (Figure 5-11), all four RCPs project quite similar changes on average (model-average warming – the black horizontal line on the bars – is within about 0.5°C). The models for RCP 8.5 have the greatest spread, particularly in summer. However, the models all agree on the direction of change (i.e. warming). For 2090 (Figure 5-12), the model spread is much larger, with the models for summer for RCP 8.5 spread across almost 4°C of warming (from ~2.1°C to ~5.9°C). All of the models agree on the direction of change (i.e. warming), aside from the lowest model under RCP 2.6 for summer, which projects no temperature change (~0°C) by 2090.




Figure 5-11: Projected seasonal temperature changes by 2040 (2031-2050) averaged over the Horizons Region, for the four RCPs. The vertical coloured bars show the range over all climate models used, and coloured stars the projected changes for each model individually (results from statistical downscaling). The black stars represent the 6-model dynamical downscaling changes. The short horizontal line is the model-average warming over all statistical and dynamical models. Blue = RCP 2.6, 23 models; green = RCP 4.5, 37 models; yellow = RCP 6.0, 18 models; red = RCP 8.5, 41 models. © NIWA.




Figure 5-12: Projected seasonal temperature changes by 2090 (2081-2100) averaged over the Horizons Region, for the four RCPs. The vertical coloured bars show the range over all climate models used, and coloured stars the projected changes for each model individually (results from statistical downscaling). The black stars represent the 6-model dynamical downscaling changes. The short horizontal line is the model-average warming over all statistical and dynamical models. Blue = RCP 2.6, 23 models; green = RCP 4.5, 37 models; yellow = RCP 6.0, 18 models; red = RCP 8.5, 41 models. © NIWA.

Figure 5-13 and Figure 5-14 show seasonal rainfall projections from all the models individually for the Whanganui grid point only for 2040 and 2090, respectively. There is disagreement between the models as to the direction of projected rainfall changes, as identified in Table 5-3 and Table 5-4 for the different RCPs. However, for 2040 in Figure 5-13, the model-average rainfall projections are quite similar for all seasons (with summer being the most variable), even though the spread of the models under each RCP is quite large (spread across approximately -10% to +20% precipitation change). For 2090 (Figure 5-14), the model spread under each RCP is much larger than in Figure 5-13 and the model-averages between each RCP are quite varied.

Note that Figure 5-13 to Figure 5-16 show the model variability at two grid points only (Whanganui and Dannevirke), rather than a regional average (as was done for temperature). This is because the projected changes to rainfall vary greatly over the region. These figures can be replicated for any grid point in the Horizons Region, upon request.




Figure 5-13: Projected seasonal rainfall changes by 2040 (2081-2100) for Whanganui, for the four RCPs. The vertical coloured bars show the range over all climate models used, and coloured stars the projected changes for each model individually (results from statistical downscaling). The black stars represent the 6-model dynamical downscaling changes. The short horizontal line is the model-average rainfall over all statistical and dynamical models. Blue = RCP 2.6, 23 models; green = RCP 4.5, 37 models; yellow = RCP 6.0, 18 models; red = RCP 8.5, 41 models. © NIWA.




Figure 5-14: Projected seasonal rainfall changes by 2090 (2081-2100) for Whanganui, for the four RCPs. The vertical coloured bars show the range over all climate models used, and coloured stars the projected changes for each model individually (results from statistical downscaling). The black stars represent the 6-model dynamical downscaling changes. The short horizontal line is the model-average rainfall over all statistical and dynamical models. Blue = RCP 2.6, 23 models; green = RCP 4.5, 37 models; yellow = RCP 6.0, 18 models; red = RCP 8.5, 41 models. © NIWA.

Figure 5-15 and Figure 5-16 show seasonal rainfall projections from all the models individually for the Dannevirke grid point only for 2040 and 2090, respectively. Although most of the ensemble-averages show an increase in precipitation (the only decreased precipitation projection is for RCP 8.5 in autumn at 2090), there is some disagreement between the individual models as to the direction and magnitude of projected rainfall changes – the spread within each RCP coloured bar is quite large, for example, ranging from a 20% decrease to more than a 40% increase in rainfall during winter for under RCP8.5 at 2090 (Figure 5-16). The model spread is larger for Dannevirke than for Whanganui for all RCPs at both 2040 and 2090.




Figure 5-15: Projected seasonal rainfall changes by 2040 (2081-2100) for Dannevirke, for the four RCPs. The vertical coloured bars show the range over all climate models used, and coloured stars the projected changes for each model individually (results from statistical downscaling). The black stars represent the 6-model dynamical downscaling changes. The short horizontal line is the model-average rainfall over all statistical and dynamical models Blue = RCP 2.6, 23 models; green = RCP 4.5, 37 models; yellow = RCP 6.0, 18 models; red = RCP 8.5, 41 models. © NIWA.




Figure 5-16: Projected seasonal rainfall changes by 2090 (2081-2100) for Dannevirke for the four RCPs. The vertical coloured bars show the range over all climate models used, and coloured stars the projected changes for each model individually (results from statistical downscaling). The black stars represent the 6-model dynamical downscaling changes. The short horizontal line is the model-average rainfall over all statistical and dynamical models. Blue = RCP 2.6, 23 models; green = RCP 4.5, 37 models; yellow = RCP 6.0, 18 models; red = RCP 8.5, 41 models. © NIWA.

5.5 Scenarios for Changes in Extreme Rainfall

A warmer atmosphere can hold more moisture (about 8% more for every 1°C increase in temperature), so there is potential for heavier extreme rainfall with global increases in temperatures under climate change. In its Fifth Assessment Report, the IPCC concluded that the frequency of heavy precipitation events is “very likely” to increase over most mid-latitude land areas (this includes New Zealand) (IPCC, 2013, Table SPM.1). Given the mountainous nature of New Zealand, spatial patterns of changes in rainfall extremes are expected to depend on changes in atmospheric circulation and storm tracks.

NIWA produced guidance on changes in heavy rainfall to be used for “screening assessments”⁴ in New Zealand, for the 2008 update to the Local Government Guidance manual (Ministry for the Environment, 2008a). The manual recommends use of a geographically uniform relationship between projected changes in temperature and changes in extreme rainfall return period statistics. An overview of the process for producing heavy rainfall statistics for screening analyses, with a detailed example of its application for Palmerston North, is provided here. This method uses augmentation amounts for various rainfall return intervals and durations set out in Table 5-5, which is a reproduction of Table 5.2 of the revised Guidance Manual (Ministry for the Environment, 2008a). The recommendation in the Local Government Guidance manual is that if a screening analysis using statistics produced through this process indicates changes in heavy rainfall could lead to problems for a particular asset or activity, then further guidance should be sought from a science provider for a more detailed risk analysis.

⁴ “Screening” describes an initial assessment step to consider whether potential impacts of climate change on a particular function or item of infrastructure are likely to be material.

Rainfall depth-duration-frequency statistics for Palmerston North under current conditions are provided in Table 5-6. Statistics for screening studies under mid-range and high-end temperature scenarios for 2100 are provided in Table 5-7 to Table 5-9.

Table 5-5: Augmentation factors (percentage increases per degree of warming) used in deriving changes in extreme rainfall for preliminary scenario studies. [Note: In preparing this table, all reasonable skill and care was exercised, using best available methods and data. Nevertheless, NIWA does not accept any liability, whether direct, indirect, or consequential, arising out of its use].

Duration	ARI						
	2 yrs	5 yrs	10 yrs	20 yrs	30 yrs	50 yrs	100 yrs
< 10 minutes	8.0	8.0	8.0	8.0	8.0	8.0	8.0
10 minutes	8.0	8.0	8.0	8.0	8.0	8.0	8.0
30 minutes	7.2	7.4	7.6	7.8	8.0	8.0	8.0
60 minutes	6.7	7.1	7.4	7.7	8.0	8.0	8.0
2 hours	6.2	6.7	7.2	7.6	8.0	8.0	8.0
3 hours	5.9	6.5	7.0	7.5	8.0	8.0	8.0
6 hours	5.3	6.1	6.8	7.4	8.0	8.0	8.0
12 hours	4.8	5.8	6.5	7.3	8.0	8.0	8.0
24 hours	4.3	5.4	6.3	7.2	8.0	8.0	8.0
48 hours	3.8	5.0	6.1	7.1	7.8	8.0	8.0
72 hours	3.5	4.8	5.9	7.0	7.7	8.0	8.0

Table 5-6: Current rainfall depth-duration-frequency statistics for Palmerston North from HIRDS V3. Numbers in the body of the table are in mm.

ARI (years)	Duration									
	10m	20m	30m	60m	2h	6h	12h	24h	48h	72h
2	6.9	9.5	11.4	15.7	20.8	32.5	43.0	57.0	68.4	76.2
5	9.1	12.6	15.2	20.9	27.2	41.3	53.8	70.1	84.2	93.8
10	11.0	15.2	18.3	25.2	32.5	48.5	62.4	80.4	96.6	107.6
20	13.2	18.2	21.9	30.2	38.4	56.5	72.0	91.7	110.2	122.7
30	14.7	20.2	24.3	33.5	42.4	61.7	78.1	99.0	118.9	132.4
50	16.7	23.0	27.7	38.1	47.9	68.8	86.5	108.8	130.7	145.5
100	19.9	27.4	33.0	45.4	56.4	79.8	99.3	123.6	148.5	165.3

Table 5-7: Projected rainfall depth-duration-frequency statistics for Palmerston North in 2100, for a low-range temperature scenario (1°C warming), from HIRDS V3.

ARI (years)	Duration									
	10m	20m	30m	60m	2h	6h	12h	24h	48h	72h
2	7.5	10.2	12.2	16.8	22.1	34.2	45.1	59.5	71.0	78.9
5	9.8	13.6	16.3	22.4	29.0	43.8	56.9	73.9	88.4	98.3
10	11.9	16.4	19.7	27.1	34.8	51.8	66.5	85.5	102.5	113.9
20	14.3	19.7	23.6	32.5	41.3	60.7	77.3	98.3	118.0	131.3
30	15.9	21.8	26.2	36.2	45.8	66.6	84.3	106.9	128.2	142.6
50	18.0	24.8	29.9	41.1	51.7	74.3	93.4	117.5	141.2	157.1
100	21.5	29.6	35.6	49.0	60.9	86.2	107.2	133.5	160.4	178.5

Table 5-8: Projected rainfall depth-duration-frequency statistics for Palmerston North in 2100 for a mid-range temperature scenario (2°C warming), from HIRDS V3.

ARI (years)	Duration									
	10m	20m	30m	60m	2h	6h	12h	24h	48h	72h
2	8.0	11.0	13.0	17.8	23.4	35.9	47.1	61.9	73.6	81.5
5	10.6	14.5	17.4	23.9	30.8	46.3	60.0	77.7	92.6	102.8
10	12.8	17.6	21.1	28.9	37.2	55.1	70.5	90.5	108.4	120.3
20	15.3	21.1	25.3	34.9	44.2	64.9	82.5	104.9	125.8	139.9
30	17.1	23.4	28.2	38.9	49.2	71.6	90.6	114.8	137.4	152.8
50	19.4	26.7	32.1	44.2	55.6	79.8	100.3	126.2	151.6	168.8
100	23.1	31.8	38.3	52.7	65.4	92.6	115.2	143.4	172.3	191.7

Table 5-9: Projected rainfall depth-duration-frequency statistics for Palmerston North in 2100, for a higher-end temperature scenario (3°C warming), from HIRDS V3.

ARI (years)	Duration									
	10m	20m	30m	60m	2h	6h	12h	24h	48h	72h
2	8.6	11.7	13.9	18.9	24.7	37.7	49.2	64.4	76.2	84.2
5	11.3	15.5	18.6	25.4	32.7	48.9	63.2	81.5	96.8	107.3
10	13.6	18.8	22.5	30.8	39.5	58.4	74.6	95.6	114.3	126.6
20	16.4	22.6	27.0	37.2	47.2	69.0	87.8	111.5	133.7	148.5
30	18.2	25.0	30.1	41.5	52.6	76.5	96.8	122.8	146.7	163.0
50	20.7	28.5	34.3	47.2	59.4	85.3	107.3	134.9	162.1	180.4
100	24.7	34.0	40.9	56.3	69.9	99.0	123.1	153.3	184.1	205.0

Projected rainfall depth-duration-frequency tables for other locations in Horizons Region can be produced using HIRDS (High Intensity Rainfall System) software package (www.hirds.niwa.co.nz) and the process is described in the revised Local Government Guidance Manual (Ministry for the Environment, 2008a). Note that a significant update to HIRDS is currently underway (as at mid 2016).

Mullan et al. (2016) calculated projected changes in extreme precipitation using the same models as in Section 5.3. The frequency of extreme precipitation, as quantified by the changes in the 99th percentile of the daily precipitation distribution (i.e. the top 1% of rain days), shows a systematic increase in much of the South Island, with both time and increasing greenhouse gas concentration (different RCPs used) (Figure 5-17). Over the North Island, however, projected changes are small and erratic. To some degree, this is because a 20-year period is too short to obtain a robust signal in the precipitation extremes. However, this preliminary analysis at least suggests that some regional variation may be expected in future return periods of extreme rainfall.

Focussing on the 2090 changes under RCP 8.5 (the strongest forcing), the only coherent regions to show a decrease in daily extreme rainfall are Northland and parts of the Wairarapa and Hawke's Bay on the east coast of the North Island (including the eastern portion of Horizons Region). These areas also show the largest reductions in winter precipitation (Figure 5-10). As a cautionary aside, the climate models being used do not have the resolution to realistically simulate tropical cyclones, and thus extreme rainfall from these phenomena are likely to be underestimated in our results.




Figure 5-17: Change in the magnitude of the 99th percentile of daily precipitation (in %). For all four RCPs and three future time periods, relative to the daily 99th percentile in the baseline 1986–2005 period. The red colours indicate a decrease in extreme rainfall, orange little change, and green, blue, and purple a progressive increase in extremes. Results are based on a 6 model ensemble average of dynamically downscaled data. After Mullan et al. (2016).

5.6 Evaporation, Soil Moisture and Drought

The increase in frequency and intensity of droughts in a changing climate is of deep concern for the New Zealand society and economy, not the least for the stakeholders of the primary sector. Drought intensity is affected by increasing temperature which in turn increases moisture loss through higher

evapotranspiration rates, and also by the lack of sufficient moderate intensity precipitation required to recharge aquifers and replenish soil moisture.

Potential evapotranspiration deficit (PED) is the cumulative sum of potential evapotranspiration (PET) from 1 July of a calendar year to 30 June of the next year, for days of soil moisture under half of available water capacity (AWC), where an AWC of 150mm for silty-loamy soils is consistent with estimates in previous studies (e.g. Mullan et al., 2005). PED, in units of mm, can be thought of as the amount of rainfall needed in order to keep pastures growing at optimum levels. As a rule of thumb, an increase in PED of 30 mm or more corresponds to an extra week of reduced grass growth.

A regional map of projected changes in potential evapotranspiration deficit is presented in Figure 5-18. The maps are plotted with a range of up to 200mm of accumulated PED anomaly with respect to the historical average.




Figure 5-18: RCM-projected changes in Potential Evapotranspiration Deficit (PED, in mm accumulation over the July-June ‘water year’) for Horizons Region. With respect to the baseline 1995 period, for RCP 4.5 and RCP 8.5, for 2040 and 2090. Black contour lines are every 40mm of PED/year. © NIWA.

Horizons Region has been separated into four sub-regions for the purposes of analysing changes to PED (and corresponding potential drought risk).

Northern Horizons sub-region (Taumarunui to Raetihi)

At 2040 under both RCP 4.5 and RCP 8.5, smaller increases in PED (compared to the rest of the region) are projected for the northern part of Horizons Region (up to 60 mm/yr). At 2090 under RCP 8.5 the increases in PED are slightly higher (up to 80 mm/yr in the north of the sub-region).

Central Horizons sub-region (Waiouru to Feilding)

At 2040 under RCP 4.5, increases in PED of up to 100 mm per year are projected in small parts of the sub-region near Taihape. By 2090 under RCP 4.5, most of the central sub-region may experience increases of 80-100 mm per year with a pocket east of Taihape projecting increases of over 120 mm per year. However, under RCP 8.5, the projections of increases in PED are more significant, with similar changes to RCP 4.5 projected for the central sub-region by 2040, but much of the sub-region projecting over 160 mm/yr, with small parts projecting a 200 mm increase per year by 2090, the highest increase in the Horizons Region.

Western coastal Horizons sub-region (Whanganui to Levin)

At 2040, the western coastal sub-region experiences increases in PED of up to 60 mm/yr under both RCP 4.5 and RCP 8.5. At 2090 under RCP 4.5, the north of the sub-region experiences increases in PED of over 80 mm/yr, while the southern part of the sub-region experiences increases of around 60 mm/yr. At 2090 under RCP 8.5, the northern part of the sub-region experiences increases of around 100 mm/yr, and the southern part of the sub-region experiences increases in PED of around 80 mm/yr.

Eastern Horizons sub-region (east of the Ruahine and Tararua Ranges)

PED has similar projections at 2040 under both RCP 4.5 and RCP 8.5 for the sub-region east of the Ruahine and Tararua Ranges – most of this sub-region may experience an increase of 80-100 mm/yr. However, by 2090, the RCPs show quite different projections, with RCP 4.5 projecting an increase of 80-100 mm/yr and RCP 8.5 projecting a 120-140 mm increase in PED per year.

Overall, the central sub-region and eastern sub-region are most at risk of future drought due to the larger projected increases in PED.

A NIWA study published in 2011 (Clark et al., 2011) used downscaled climate model results from the IPCC Fourth Assessment Report to examine how the frequency of very dry conditions could change over the 21st century. Three major global greenhouse gas emissions scenarios were used (B1, A1B, and A2), and the final estimates of drought probability were derived from a nationally comprehensive soil moisture indicator.

The study established distinct regional differences across New Zealand in changes to drought vulnerability projected under future climate change, with an increase in drought on the east coast of the North and South Islands being the most plausible and consistent outcome. This is consistent with previous studies on climate change impacts on drought in a New Zealand context (e.g. Mullan et al., 2005). The study concluded that drought risk is expected to increase during this century in all areas that are currently drought prone, under both the ‘low-medium’ and ‘medium-high’ scenarios. The ‘drought risk’ was analysed in terms of soil moisture levels – drought initiation occurs when soil moisture falls below the historically established 10th percentile for the given time of year for a period greater than one month, and drought termination occurs when soil moisture is above the 10th percentile for one month.

Under the most likely mid-range emissions scenario the projected increase in percentage of time spent in drought from 1980-99 levels is about 5% for 2030-2050 and 10% for 2070-2090 for the Horizons Region. This can be interpreted as: for a site that is currently in drought 5% of the time in the Manawatu Plains, in 2030-2050 it is likely that this same location will be in drought 10% of the time (i.e. an additional 5%), and in 2070-2090 that location is likely to be in drought 15% of the time (an additional 10%). The Central Plateau is less likely to be affected as the remainder of the region by climate-change-induced drought.

Mullan et al. (2016) calculated the number of dry days (days with rainfall <1mm) for different RCPs (Figure 5-19). Blue and green shading indicates a decrease in dry days (i.e., more rain-days), yellow little change, and orange and red an increase in the number of dry days per year. The frequency of dry days increases with time and RCP for much of the North Island, and for high altitude inland regions in the South Island. The frequency of dry days decreases on the west and east coastal regions in the South Island.




Figure 5-19: RCM-projected changes in the annual number (in days) of “dry days” (precipitation below 1 mm/day). With respect to the baseline 1995 period, for all four RCPs and three future time periods. Blue shades indicate a decrease in dry days and red shades indicate an increase in dry days. Results are based on a 6 model ensemble average of dynamically downscaled data. After Mullan et al. (2016).

5.7 Pressure and Wind

Mean sea-level pressure and wind projections have been derived from the Regional Climate Model (RCM) simulations by Mullan et al. (2016). The RCM simulations provide much more information about weather parameters than is readily available from statistical downscaling. In all cases, there is a maximum of six models available for analysis for each RCP and time period; four models are available beyond 2100 for RCPs 2.6 and 8.5, five models for RCP 4.5, and only one model for RCP 6.0.

The key projected changes in mean sea-level pressure (MSLP) and mean winds are as follows (more detail can be found in Mullan et al. (2016)):

- MSLP tends to increase in summer (December-January-February, DJF), especially to the south-east of New Zealand. In other words, the airflow becomes more north-easterly, and at the same time more anticyclonic (high pressure systems).
- MSLP tends to decrease in winter (June-July-August, JJA), especially over and south of the South Island, resulting in stronger westerlies over central New Zealand.
- In the other seasons (autumn and spring), the pattern of MSLP change is less consistent with increasing time and increasing emissions. However, there is still general agreement for autumn changes to be similar to those of summer (i.e., more anticyclonic), and for spring changes to be similar to those of winter (lower pressures south of the South Island, and stronger mean westerly winds over southern parts of the country).

The 99th percentile of daily-mean wind speed was evaluated over the historical 1986-2005 period at each VCSN grid-point in the downscaled (but not bias-corrected) regional model output data, by Mullan et al. (2016). Figure 5-20 maps how the 99th percentiles at future periods differs from the current climate for each of the four RCPs.

In Figure 5-20, yellow shading means little or no change from present, green a decrease in extreme wind speed, and red an increase. For most of the RCPs and time periods, the southern half of the North Island and all the South Island are shown as having stronger extreme daily winds in future. This is especially noticeable in the South Island east of the Southern Alps. The regional model is able to resolve speed-up in the lee of the mountain ranges, and shows increases of up to 10% or greater in Marlborough and Canterbury by the end of the century under the highest RCP8.5 forcing. However, there is a decrease in extreme winds in the North Island from Northland to Bay of Plenty, probably because of increasing anticyclonic conditions.

No seasonal breakdown of extremes is given, but it is expected that the higher winds in the east of the South Island are primarily due to the increased westerly pressure gradient in winter and spring. Very localised extreme winds from more vigorous summer convection are also potentially a problem in the future, but such events are not resolved by the regional model being used here.




Figure 5-20: Change in the magnitude of the 99th percentile of daily-mean wind speed, for all four RCPs and three future time periods. Relative to the daily 99th percentile in the baseline 1986–2005 period. Blue and green shades indicate a decrease in wind speed and red shades indicate an increase in wind speed. Results are based on a 6 model ensemble average of dynamically downscaled data. After Mullan et al. (2016).

5.8 Hill Country Erosion

Hill country erosion is a significant issue for Horizons Region, due to the high portion of the region that is in hill country landscapes, and the importance of pasture-based agriculture to the region. Major erosion events in the past have caused significant economic stress for the region.

Basher et al. (2012) studied the impacts of climate change on erosion. They concluded that the main features of climate change that will affect erosion are:

- Changes in rainfall patterns (increasing annual rainfall in the west of the Horizons Region and decreasing in the east, as shown in Section 5.3) and extreme rainfall (Section 5.4)
- Increases in temperature affecting plant water use and soil water balance (see Section 5.1 and 5.5)
- Increased windiness and incidence of drought (see Section 5.5 and 5.6)

Climate and erosion are clearly linked through water movement into and through the soil, the soil water balance, and slope hydrology. Projected increases in extreme rainfalls in some areas (Section 5.4) will play a critical role in determining the effect of climate change. Hillslope erosion processes (e.g. shallow landslides, earthflows, gully, and sheet erosion) are likely to be influenced by climate change.

Shallow rapid landslides are usually triggered by a rainfall event, and as such they are likely to change in frequency with climate change depending on changes to extreme and annual rainfall, rainfall variability, ex- and extra-tropical cyclone variability, and wind.

Earthflow erosion (extensive on the soft rock hill country of the northern part of the Horizons Region) has complex links to soil moisture and storm rainfall. Changes in rainfall and temperature (through its effect on evapotranspiration) with climate change may influence rates of earthflow movement.

Gully erosion and sheet erosion is related to high annual and storm rainfalls, so any increase in rainfall with climate change (either of annual totals or extreme events) can be expected to increase these types of erosion.

Some parts of Horizons Region were identified by Basher et al. (2012) as being more susceptible to erosion with the effects of climate change:

- Areas most susceptible to increased landsliding include: the soft rock hill country of Taranaki, southern Waikato, Horizons Region west of the Ruahine Range (the regions outside of Horizons have been included here as they border on Horizons Region).
- Areas most susceptible to increased gully erosion include: the soft rock hill country of inland Taranaki, and Whanganui areas. Most of the areas with the highest potential for earthflow erosion (including the eastern portion of Horizons Region, east of the Ruahine and Tararua Ranges) are projected to have a decrease in rainfall, which is likely to reduce rates of earthflow movement.
- Areas most susceptible to increased sheet erosion are the Volcanic Plateau (including the intensive cropping area around Ohakune), and the northern Horizons-Taranaki hill country.

As stated in Section 5.4, there are small and erratic daily extreme precipitation changes in Horizons Region (Mullan et al., 2016), so it is unclear whether daily extreme precipitation changes will have an impact on hill country erosion in the region. However, the seasonal precipitation projections in Section 5.3 suggest that precipitation may increase for the western part of Horizons Region (and especially the northwest part of the region in winter for RCP 8.5 at 2090 (Figure 5-10)). This part of Horizons Region was identified by Basher et al. (2012) as being particularly susceptible to erosion.

5.8.1 Climate change impacts on sediment loads in rivers

A report examined the implications of climate change on sediment loading in rivers in Horizons Region, considering the management scenarios of the Sustainable Land Use Initiative (SLUI) (Manderson et al., 2015). The report modelled the effects of climate change on sediment yields for rivers, assuming that SLUI continues according to management scenario #3. Climate change projections are based on the IPCC Fourth Assessment Report.

In this case, ‘climate change’ was considered to be the change in extreme storm rainfall driven by temperature increases. Three different climate change scenarios were modelled, which used the SRES emissions scenarios from the IPCC Fourth Assessment Report. A ‘minor’ climate change impact scenario was modelled as halfway between the status quo and scenario A1B, a ‘medium’ scenario used the A1B levels, and a ‘major’ scenario used A1F1 levels. The sediment loading model used was SedNetNZ.

Climate change (i.e. increased storm rainfall) is projected to increase sediment loading in Horizon Region’s rivers from a baseline of 9.81 Mt/yr to 10.83 Mt/yr under a climate change with minor impact (+10.4%), 11.85 Mt/yr for the medium impact scenario (+20.8%), and 12.71 Mt/yr for the major impact scenario (+29.6%). However, climate change implications on sediment loads vary according to catchment climate and type of erosion terrain. Although all scenarios project an increase in sediment loading in rivers, these estimates are still less than 2004 pre-SLUI levels.

However, climate change will reduce the long-term effectiveness of SLUI, with lower rates of sediment reduction under management scenario #3. Adopting other management scenarios may improve long-term sediment reduction under climate change.

The reader is directed to Manderson et al. (2015) for more details about the study.

5.9 Solar Radiation

Mullan et al. (2016) produced projections of solar radiation for the four RCPs at different periods during the 21st century.

The geographic distribution of solar radiation depends not only on astronomical factors but also local rainfall (and cloudiness) patterns. Currently, the highest solar radiation levels are recorded in Nelson-Marlborough and in central Otago regions in summer months, and in northern North Island and Nelson-Marlborough in winter months. Astronomical factors will not change over the projection period but cloud will.

The seasonal patterns of projected changes are clearest for the most extreme RCP (RCP 8.5). The solar radiation projection suggests increases of up to 10% on the West Coast in summer, and smaller increases elsewhere with notable exception of the coastal Canterbury where solar radiation is predicted to decrease. Autumn solar radiation patterns are similar but weaker to those of summer. The winter changes are almost the reverse of the summer ones: about a 5% decrease in radiation in western parts of the North Island, and 10% or more in western and southern South Island. Eastern North Island is projected to have an increase in winter solar radiation levels. In spring, most of the North Island, and the northern third of the South Island, displays an increase in radiation, with decreases further south.

For more detail on solar radiation projections, see Mullan et al. (2016).

5.10 Climate Change and Sea Level

Sea levels will continue to rise over the 21st century and beyond, primarily because of thermal expansion within the oceans and loss of ice sheets and glaciers on land. The basic range of projected global sea-level rise estimated in the IPCC's Fifth Assessment Report (IPCC, 2013) is for a rise of 0.26 m to 0.82 m for 2081-2100 (2080s and 2090s) relative to the average sea level over the period 1986-2005, as shown in Figure 5-21. This is based on projections from IPCC AR5 climate model projections in combination with process-based models of glacier and ice sheet surface mass balance for the four different RCP emissions scenarios. Global mean sea level rise for the scenarios will likely be in the 5 to 95% ranges characterising the spread of the model results (bars on the right hand side of Figure 5-21).




Figure 5-21: Projections of global mean sea level rise over the 21st century relative to 1986-2005.

Projections are from the combination of the Coupled Model Intercomparison Project (CMIP5) ensemble with process-based models, for RCP2.6 and RCP8.5. The assessed likely range is shown as a shaded band. The assessed likely ranges for the mean over the period 2081-2100 for all RCP scenarios are given as coloured vertical bars, with the corresponding median value given as a horizontal line. After IPCC (2013).

In all emissions scenarios, thermal expansion is the largest contribution to global mean sea-level rise, accounting for about 30-55% of the total. Glaciers are the next largest, accounting for 15-35% of total sea level rise. By 2100, 15-55% of the present glacier volume is projected to be eliminated under the lowest emissions scenario, and 35-85% under the highest emissions scenario. The increase in surface melting in Greenland is projected to exceed the increase in accumulation, and there is high confidence that the surface mass balance changes on the Greenland ice sheet will make a positive contribution to sea-level rise over the 21st century. On the Antarctic ice sheet, surface melting is projected to remain small.

Figure 5-22 shows Wellington and Auckland annual mean sea level measurements spliced with global-mean sea-level rise projections for two of the RCPs (RCP2.6 and RCP8.5) from IPCC (2013). Note that IPCC provided a caveat that further collapse of Antarctica ice sheets could cause global sea level to rise substantially above the *likely* ranges by 2100 (shown as dashed lines in Figure 5-22), with medium confidence that the additional contribution would not exceed several decimetres of sea-level rise by 2100. The RCP projections from IPCC have been extended out from 2100 to 2120, to assist with the application of the NZ Coastal Policy Statement that requires coastal hazards and

climate change effects be assessed over “at least 100 years”. Both measurements from the two ports and the projections are relative to a baseline averaged from 1986–2005 (centred on 1996).




Figure 5-22: AR5 global mean sea-level projections extended to 2120 spliced with annual MSL from Wellington and Auckland. Dashed lines show the upper and lower bounds of the *likely* range for each RCP shown, which in the calibrated language of IPCC means there is a 33% chance the SLR could lie outside those bounds (Church et al., 2013a). Source: QA for annual MSL for historic NZ ports undertaken by Prof. John Hannah and NIWA – data sourced originally from the port companies and obtained from the Land Information NZ archives. The global-mean SLR projections were interpolated by data from Table 13.5 (Church et al., 2013b) and from Figure SPM.9 in the IPCC AR5 Summary for Policymakers (IPCC, 2013).

Sea level rise will have an impact on sediment redistribution, the partitioning of habitats within estuaries, salinity, tidal range, and submergence periods, with consequences for biological communities in the estuaries as well as human communities that exist nearby (Wong et al., 2014). Estuaries and coastal lagoons may shrink because landward migration is restricted due to human occupation, or extend due to the drowning of marshes and coastal wetlands.

A manual for local government on coastal hazards and climate change was published in 2008 (Ministry for the Environment, 2008b). This guidance manual uses projections based on the IPCC’s Fourth Assessment Report. In time, it is expected an updated report will be published using projections based on the IPCC Fifth Assessment Report projections, but in this report the 2008 report will be referenced.

The Coastal Hazards and Climate Change Guidance Report includes suggestions on changes in mean sea-level for use in future planning and decisions. Numbers for use in such guidance depend on risk management considerations as well as scientific assessment. The guidance manual advocates the use of a risk assessment process to assist incorporating sea-level rise and the associated uncertainties, within local government planning and decision-making. This requires a broader consideration of the potential impacts or consequences of sea-level rise on a specific decision or issue. Rather than define a specific climate change scenario or sea-level rise value to be accommodated, it is recommended in

the manual that the magnitude of sea-level rise accommodated is based on the acceptability of the potential risk.

To aid this risk assessment process, the manual recommends that allowance for sea-level rise is based on the IPCC Fourth Assessment Report, and that consideration be given to the potential consequences from higher sea-levels due to factors not included in current global climate models⁵.

For planning and decision timeframes out to the 2090s (2090-2099):

- a. A base value sea level rise of 0.5 m relative to the 1980-1999 average should be used, along with:
- b. An assessment of the potential consequences from a range of possible higher sea-level rises (particularly where impacts are likely to have high consequence or where additional future adaptation options are limited). At the very least, all assessments should consider the consequences of a mean sea-level rise of at least 0.8 m relative to the 1980-1999 average.

For planning and decision timeframes beyond 2100 where, as a result of the particular decision, future adaptation options will be limited, an allowance of sea-level rise of 10 mm per year beyond 2100 is recommended (in addition to the above recommendation). Climate change will also impact on other coastal hazard drivers, such as tides, storm surge, waves, swell, and coastal sediment supply. The potential changes and their impacts are at present much less well understood, but the manual provides pragmatic guidance informed by expert judgement and the current state of scientific knowledge.

A small amount of research has been completed within Horizons Region concerning the effects of sea-level rise on coastal morphology. The following reports state that sea-level rise and extreme sea levels have and will have an impact on coastal flooding and morphology in Horizons Region, at present and into the future. Note that these reports use the general sea-level rise value from (Ministry for the Environment, 2008b); no sea-level rise estimates have been generated specifically for Horizons Region. The reader is directed to the following reports for location-specific information on the degree of landward beach movement and changes to extreme sea-level elevation with climate-induced sea-level rise:

- Bell (2015) – Himatangi Beach
- Tonkin & Taylor (2010) – Akitio and Herbertville
- Tonkin & Taylor (2013) – Waitarere Beach, Hokio Beach, Waikawa Beach
- Blackwood (2007) and Shand (2008) – Foxton Beach
- Bell (2014) – Koitiata Beach and Castlecliff Beach

5.10.1 Effect of Sea-Level Rise on High Tide Exceedance Frequency

The Coastal Hazards and Climate Change Guidance Manual (Ministry for the Environment, 2008b) provides information on tide ranges and frequency of high tides. Tidal ranges and the timing of high and low tides in shallow harbours, river mouths and estuaries could be altered by changes in channel depth due to climate change. These changes could occur through either the deepening of channels

⁵ Such factors relate to uncertainties associated with increased contribution from the Greenland and Antarctica ice sheets, carbon cycle feedbacks, and possible differences in mean sea level when comparing the New Zealand region with the global average.

where sea-level rise exceeds the rate of sediment build-up, or conversely by the formation of shallower channels where rates of sediment build-up (i.e. from increased runoff due to more intense rainfall events) exceeds sea-level rise.

The present Mean High Water Spring level around New Zealand coastlines will be exceeded much more frequently by high tides in the future, particularly on sections of the coast where the tide range is relatively small (compared with those sections of the coast where the tide range is relatively large). Problems will be exacerbated for coastlines with smaller tidal ranges in proportion to sea-level rise, where high tides will more often exceed current upper-tide levels, thus allowing more opportunity to coincide with storms or large swell.

Sea-level rise will have a greater influence on storm inundation and rates of coastal erosion on the central parts of the east coast (Napier/Gisborne) and Cook Strait/Wellington areas (due to their smaller tidal range) than on coastal regions with larger tidal ranges (e.g. west coast – Taranaki, Nelson, Westport).

A report on New Zealand's coastal sensitivity by Goodhue et al. (2012a) found that the east coasts of both the North and South Islands are more sensitive to erosion and inundation caused by climate change, because of a combination of factors such as wave exposure, relatively low tidal ranges, sediment budget deficits, and proximity to tidal inlets. Conversely, west coast shores are less sensitive to climate-driven change, mainly because they are already regularly exposed to high wave energy.

An example of the effect that future sea-level rise has on the frequency of high tides at Foxton is shown in Figure 5-23 and Table 5-10. The black line in Figure 5-23 shows the present-day sea levels and the upper curves are for different levels of sea-level rise. Currently, 10% of high tides exceed the MHWS-10 level (Mean High Water Spring level that 10% of high tides exceed, excluding weather and climate effects). With 0.4 m of sea-level rise, approximately 60% of high tides will exceed this level and with 0.8 m of sea-level rise about 99% of high tides will exceed this level, again excluding weather and climate effects such as storm surge.




Figure 5-23: The frequency of occurrence of high tides exceeding different present day tide marks at Foxton. The tide marks for the present is shown by the heavy line, 0.4m of sea-level rise (blue line) and 0.8m of sea-level rise (red line). ©NIWA.

Table 5-10: Data analysis of effect of sea-level rise at Foxton on components of tide. © NIWA.

Maximum HW	1.375 m
MHWPS (Mean High Water at Perigean Spring) 1.203 m exceeded by	4.2%
Pragmatical MHWS for 10% exceedance	1.12 m
MHWS (Mean High Water Spring)	1.068 m exceeded by 15.3%
MHWN (Mean High Water Neap)	0.432 m exceeded by 92.0%
MHWAN (Mean High Water at Apogean Neap)	0.297 m exceeded by 99.2%
Minimum HW	0.196 m
Minimum SLR for ALL high tides to exceed present MHWPS	1.01 m
Minimum SLR for ALL high tides to exceed present MHWS-10%ile	0.92 m
% of high tides that would exceed current MHWS-10 for 0.4 m SLR	60.2%

% of high tides that would exceed current MHWS-10 for 0.8 m SLR	98.6%
--	-------

5.11 Climate Change Impacts on Other Coastal Hazard Drivers

While it is expected that the intensity of tropical cyclones and extratropical cyclones will increase (i.e. wind speed and rain rates), it is likely that their frequency will either decrease or remain essentially unchanged (IPCC, 2013). These storms affect the coastal zone through impacts on waves, storm surge, and swell.

Some high-resolution atmospheric models in the IPCC Fifth Assessment Report have realistically simulated tracks and counts of tropical cyclones, and models are able to capture the general characteristics of storm tracks and extratropical cyclones with evidence of improvement since the IPCC Fourth Assessment Report. However, uncertainties in projections of cyclone frequency and tracks make it difficult to project how future changes will impact particular regions.

In addition, the projections of storm surges (increase in sea level caused by the inverse barometer effect from large storms such as tropical cyclones) have low confidence, in part due to the high uncertainty surrounding future storminess. Changes in storm surge will depend on changes in the frequency, intensity, and/or tracking of low-pressure systems, and the occurrence of stronger winds associated with these systems (IPCC, 2013). Expected changes in wind and atmospheric patterns, storms and cyclones around New Zealand and the wider southwest Pacific and Southern Ocean regions also have the potential to change the wave climate experienced around New Zealand in the future. In turn, this will influence patterns of coastal erosion and the movements of beach and nearshore sediments within coastal zones.

At a large scale, it is likely that the mean significant wave heights will increase in the Southern Ocean as a result of enhanced westerly wind speeds, especially in the austral winter months (5-10% higher at the end of the 21st century than the present-day mean). In addition, Southern Ocean-generated swells are likely to affect heights, periods, and directions of waves in adjacent basins (IPCC, 2013).

A number of reports on coastal hazard assessment have been completed for Horizons Region (Bell, 2014, Bell, 2015, Blackwood, 2007, Shand, 2008, Tonkin & Taylor, 2010, Tonkin & Taylor, 2013). These reports include information on projected elevations of tide, storm surge, wave setup, and wave run up for 1% AEP events, incorporating sea-level rise projections for ~2060 and ~2110 (year of projection varies with report). These results are useful for determining coastal setback zones, but do not incorporate climate change-induced changes to waves, storm surges, and tides over and above sea-level rise. NIWA reports for other regional councils (e.g. Goodhue et al., 2012b) have used the IPCC emissions scenarios to run simulations of storm surge, storm tide height, and significant wave height under climate change to 2100.

Horizons Regional Council administers a large section of coastline on both east and west coasts, and many communities and infrastructure are placed near the coast. Therefore, more research on the effect of sea-level rise and climate change on coastal hazard drivers (e.g. storm surge and waves) could be useful for the region, in order to better understand the potential impacts of climate-induced changes for the region's coasts.

5.12 Hydrological Impacts of Climate Change

The climate projections used to drive the hydrological model TopNet in order to make future projections of hydrological changes in New Zealand rivers are from the Regional Climate Model (RCM) simulations by Mullan et al. (2016). Results are presented in Table 5-11 for the Manawatu River in Horizons Region in terms of percentage changes relative to modelled historical conditions (1986-2005) for six different global climate models, four RCPs and two projection timelines (2031-2050: '2040' and 2081-2100: '2090'). Interpretation of each variable follows the table. Due to the log-normality of streamflow generation processes, results will be presented in term of change of median rather than of change in average (as per the climate section).

Table 5-11: Projected changes in flow variables of the Manawatu River (in %) between 1986-2005 and 2031-2050 (2040) and 2081-2100 (2090). The changes are given for four RCPs, where the ensemble median is taken over six models. The values in each column represent the ensemble median, and in brackets the range (minimum to maximum) over all models within that ensemble.

Variable and RCP	2040 Median (min, max)	2090 Median (min, max)
Mean annual flow		
rcp 8.5	1 (-7, 12)	-2 (-13, 3)
rcp 6.0	3 (-8, 10)	1 (-8, 5)
rcp 4.5	2 (-3, 10)	-2 (-5, 8)
rcp 2.6	1 (-3, 11)	3 (-2, 9)
Mean annual flood		
rcp 8.5	25 (3, 35)	13 (-5, 49)
rcp 6.0	18 (0, 29)	18 (-26, 76)
rcp 4.5	14 (4, 28)	8 (-20, 69)
rcp 2.6	26 (-15, 95)	8 (-13, 45)
Mean annual low flow (Q10)		
rcp 8.5	-7 (-17, 22)	-12 (-25, 5)
rcp 6.0	-1 (-19, 7)	-7 (-16, 8)
rcp 4.5	-5 (-12, 14)	-4 (-16, 7)
rcp 2.6	-1 (-10, 12)	1 (-15, 14)
Average seasonal flow: DJF		
rcp 8.5	0 (-19, 15)	-14 (-24, 7)
rcp 6.0	0 (-14, 24)	3 (-34, 8)
rcp 4.5	-7 (-12, 28)	3 (-12, 16)
rcp 2.6	-1 (-9, 27)	-4 (-19, 23)
Average seasonal flow: MAM		
rcp 8.5	3 (-11, 6)	-6 (-18, 2)
rcp 6.0	4 (-10, 9)	-4 (-13, 10)
rcp 4.5	2 (-12, 12)	-7 (-11, 10)
rcp 2.6	0 (-5, 16)	1 (-12, 11)
Average seasonal flow: JJA		
rcp 8.5	0 (-5, 13)	5 (-7, 9)

	rcp 6.0	2 (-10, 7)	3 (-3, 8)
	rcp 4.5	1 (0, 5)	-2 (-4, 9)
	rcp 2.6	6 (-8, 12)	3 (0, 5)
Average seasonal flow: SON			
	rcp 8.5	-1 (-10, 7)	-8 (-12, 9)
	rcp 6.0	2 (-6, 13)	-2 (-13, 7)
	rcp 4.5	8 (-4, 10)	-6 (-8, 11)
	rcp 2.6	5 (-3, 21)	3 (-4, 10)
FRE3: Number of hours above FRE3			
	rcp 8.5	2 (-23, 55)	21 (-4, 71)
	rcp 6.0	-1 (-20, 19)	23 (-6, 35)
	rcp 4.5	-5 (-16, 46)	13 (-14, 35)
	rcp 2.6	-1 (-18, 24)	0 (-16, 34)

Mean annual flow: Mean river flows change in response to shifts in precipitation patterns. The median result across all climate models of the mean flow changes at both 2040 and 2090, for all RCPs, are generally negligible (within $\pm 5\%$) with a relatively small ensemble spread, in other words projecting minimal change in mean annual flow to the end of the 21st century for the Manawatu River (Table 5-11).

Mean annual flood: The mean annual flood is the average of the maximum flood discharges experienced in a particular river, which should have a recurrence interval of once every 2.33 years. Table 5-11 shows large (>10%) increases in mean annual flood for the median ensemble result across most RCPs at both time periods, with larger increases at 2040 than at 2090 (except for RCP 6.0 ensemble median which remains the same). Note that the spread of the ensemble is quite large and increases with time.

Mean annual low flow (Q10): The mean annual low flow (taken as Q10) is defined as the flow that is exceeded 90% of the time, independently of recurrence. The median ensemble results show projected decreases in mean annual low flow for all RCPs at both time periods except for RCP 2.6 at 2090 (median = 1% increase). The largest median ensemble change is projected for 2090 under RCP 8.5, which is a reduction in mean annual low flow of 12%. However, the range of model results is quite large.

Average seasonal flow: Examining average seasonal flows shows a more nuanced picture of changing flow patterns than at the annual time scale. For many RCPs at both time periods, the median model result is within $\pm 5\%$ change, and the range of model projections is quite large. However, under RCP 8.5 at 2090 for summer, flows are projected to decrease by 14% (median model result). This is consistent with small projected decreases in summer precipitation for the area around the Manawatu River for the same RCP and time period. Spring flows are projected to slightly increase to mid-century (2040) but decrease by late-century (2090) for most RCPs.

Number of hours above FRE3: FRE3 is described by Booker (2013) as the average number of high flow events (floods or floods) per year that exceed three times the median flow. FRE3 has been used as an index relating to flow-driven disturbance in ecological studies of in-stream biota. A decrease (increase) in FRE3 would mean fewer (more) floods capable of disturbing aquatic ecosystems, suggesting an increase (decrease) in periphyton cover. Table 5-11 shows small projected

decreases in the median model result for FRE3 at 2040 under all RCPs except RCP 8.5 (2% increase), and large increases (>10%) in FRE3 at 2090 under all RCPs except RCP 2.6 (0% change).

The following caveats should be noted alongside the river flow projections presented in Table 5-11:

- All results have been reported as percent changes. There has been no attempt made by NIWA to bias correct or otherwise adjust the TopNet model output, based on observed flows.
- Low flows are taken to be the flow that is exceeded 90% of the time. There were no attempts in this analysis to define change in terms of the 7 day Mean Annual Low Flow.
- FRE3 are reported as the occurrence of discharge above three times the median flow. As per the low flow characteristics, there was no attempt made in this analysis to correctly define a FRE3 event.
- The changes are reported in terms of the change to the median of an ensemble of six RCM runs. The maximum and minimum range of change are provided across all RCPs only as a potential indication of the range of change.

5.13 Ocean Acidification

Since the beginning of the industrial era, the pH of global ocean surface water has decreased by 0.1, corresponding to a 26% increase in hydrogen ion concentration (IPCC, 2013). It is virtually certain that the increased storage of carbon by the ocean will increase acidification in the future, continuing the observed trends of the past decades. Ocean acidification in the surface ocean will follow atmospheric CO₂ and it will also increase in the deep ocean as CO₂ continues to penetrate the abyss (Figure 5-24).

A global increase in ocean acidification is projected under all RCP scenarios, due to the increasing uptake of carbon by the ocean. The corresponding decrease in surface ocean pH by the end of the 21st century is in the range of 0.06-0.07 for RCP 2.6, 0.14-0.15 for RCP 4.5, 0.20-0.21 for RCP 6.0, and 0.30-0.32 for RCP 8.5.




Figure 5-24: Change in ocean surface pH. Time series (model averages and minimum to maximum ranges and (b) maps of multi-model surface ocean pH for the scenarios RCP 2.6, RCP 4.5, RCP 6.0 and RCP 8.5 in 2081-2100. The maps in (b) show change in global ocean surface pH in 2081-2100 relative to 1986-2005. The number of CMIP5 models to calculate the multi-model mean is indicated in the upper right corner of each panel. Figure after (IPCC, 2013).

Figure 5-25 shows the projected pH decrease to 2100, using RCP 8.5, in New Zealand's Exclusive Economic Zone (EEZ). The pH decrease, from current values of ~8.08 to 7.95 by mid-century and 7.75 by 2100, is consistent with global trends of a decline by 0.3-0.4 by the end of the century. The sinusoidal pattern reflects the seasonal shift within each year of higher pH in summer (when phytoplankton growth removes CO₂) and lower pH in winter (when growth is low and mixing raises surface water CO₂).




Figure 5-25: Projected mean surface pH for the NZ EEZ open ocean from a suite of six CMIP5 models, using the RCP 8.5 scenario. The outputs from the models that show the closest fit to carbonate observations in the NZ EEZ during the present period are highlighted in red (S. Mikaloff-Fletcher, NIWA).

The 15-year Munida time-series in Figure 5-26 shows that ocean acidification of NZ waters is already evident, with an increase in dissolved surface CO₂ and associated decreases in surface pH and carbonate saturation state. The increase in dissolved CO₂ is consistent with the regional increase in atmospheric CO₂ recorded at the NIWA Baring Head Atmospheric Station. The observed decline in pH and carbonate saturation are consistent with observations at 6 other time-series stations in the global ocean, although the rate of change of pH at the Munida station is the lowest.

The variability and rate of change in pH will differ in coastal waters as these are also influenced by terrestrial factors and run-off. The rate and magnitude of acidification in coastal waters is being monitored by the recently initiated New Zealand Ocean Acidification Observing Network (NZOA-ON) of 14 stations around the coast.




Figure 5-26: Time series of Surface seawater pCO₂ (μatm , top panel), pH (middle panel), and saturation state of the carbonate mineral, aragonite (lower panel). This time series is from Sub-Antarctica water at the

Munida site (Otago shelf). Coloured symbols are the anomalies and the grey symbols the observed data, with the annual trends (yr-1) shown (Bates et al., 2014).

5.13.1 New Zealand-specific Impacts of Ocean Acidification

- There is evidence of an increase in bacterial enzyme activity under increased dissolved CO₂, which increase oxygen removal and decrease carbon uptake in the ocean (Burrell, 2015, Maas et al., 2013).
- Time series studies show no discernible impact of current increases in dissolved carbon dioxide on phytoplankton or zooplankton with carbonate shells in NZ waters currently (S. Nodder & C. Law, pers. comm.), although decreased carbonate production under conditions projected for 2100 in comparative species from other regions suggest they may be detrimentally impacted by the end of the century.
- Nitrogen fixers that live in nutrient-depleted regions are predicted to be “winners” from ocean acidification; however no significant effect of increased dissolved CO₂ was observed in experiments carried out on mixed plankton communities in NZ subtropical waters (Law et al., 2012).
- Macroalgae community structure in coastal regions may be altered in response to ocean acidification with a decline in encrusting coralline algae that use carbonate, while red algae found in deeper waters may benefit from an increase in dissolved CO₂ (Hepburn et al., 2011, Tait, 2014). Changes in the biomineralisation or species distribution of Coralline red algae may occur in response to ocean acidification, particularly in species producing high-Mg calcite (James et al., 2014, Smith et al., 2013).
- Sponges may benefit from ocean acidification (Bell et al., 2013) although those that produce calcite of high magnesium content, or aragonite may be vulnerable to dissolution (Smith et al., 2013).
- The projected decrease of carbonate saturation in the deep ocean may cause a decline in the abundance and distribution of cold water corals, which support important ecosystems in regions such as the Chatham Rise (Bostock et al., 2015). It is suggested that seamounts and topographic features may be important future refugia for cold water corals (Thresher, 2015, Tittensor et al., 2010).
- There is clear evidence of malformation of Sea Urchin larvae, in tropical to Antarctic species including from NZ, under higher dissolved CO₂. This may result in smaller larvae and an increased duration in the planktonic phase, reducing the chances of survival to the adult stage (Byrne et al., 2013, Clark et al., 2009).
- Experimental work on the impacts of acidification in New Zealand waters on juvenile paua has shown that while survival was not affected, growth was significantly reduced, and dissolution of the shell surface was evident (Cunningham, 2013). Similar effects were found for growth and shell surfaces of flat oysters (Cummings et al., 2013, Cummings et al., 2015). This is consistent with observed negative effects of ocean acidification on the function and metabolism of Antarctic bivalves (Bylenga et al., 2015, Cummings et al., 2011).

- The behaviour of Australian reef fish is affected by ocean acidification, with olfaction, hearing, visual risk assessment and activity altered due to the impact on neurotransmitter function (Munday et al., 2014). MPI funding has supported studies of the impacts of ocean acidification on Kingfish, and this work will be extended to Snapper.

5.14 Considering both Anthropogenic and Natural Changes

Much of the material in Sections 5.1 to 5.13 focuses on the projected impact on the climate of Horizons Region over the coming century of increases in global anthropogenic greenhouse gas concentrations. But natural variations, such as those described in Section 4.3 (associated with for example El Niño, La Niña, the Interdecadal Pacific Oscillation, the Southern Annular Mode, and “climate noise”), will also continue to occur. As noted at the beginning of Section 5, those involved in (or planning for) climate-sensitive activities in the Horizons Region will need to cope with the sum of both anthropogenic change and natural variability.

An example of this for temperature (from an overall New Zealand perspective) is shown in Figure 5-27. This figure shows annual temperature anomalies relative to the 1986-2005 base period used throughout this report. The solid black line on the left-hand side represents NIWA’s 7-station temperature anomalies (i.e., the average over Auckland, Masterton, Wellington, Nelson, Hokitika, Lincoln, and Dunedin), and the dashed black line represents the 1909-2014 trend of 0.92°C/century extrapolated to 2100. All the other line plots and shading refer to the air temperature averaged over the region 33-48°S, 160-190°W, and thus encompasses air temperature over the surrounding seas as well as land air temperatures over New Zealand. Post-2014, the two line plots show the annual temperature changes (for the ‘box’ average) under RCP 8.5 (orange) and RCP 2.6 (blue); a single model (the Japanese ‘*miroc5*’ model, see Mullan et al. (2016)) is selected to illustrate the interannual variability. (Note that a single illustrative model (*miroc5*) has been used in Figure 5-27 rather than the model-ensemble, which would suppress most of the interannual variability). The shading shows the range across all AR5 models for both historical (41 models) and future periods (23 for RCP 2.6, 41 for RCP 8.5).

Over the 1909-2014 historical period, the 7-station curve lies within the 41-model ensemble, in spite of the model temperatures including air temperature over the sea, which is expected to warm somewhat slower than over land (Mullan et al., 2016). For the future 2015-2100 period, the RCP 2.6 ensemble shows very little warming trend after about 2030, whereas the RCP 8.5 ensemble ‘takes off’ to be anywhere between +2°C and +5°C by 2100. The *miroc5* model is deliberately chosen to sit in the middle of the ensemble, and illustrates well how interannual variability dominates in individual years: the *miroc5* model under RCP 8.5 is the warmest of all models in the year 2036 and the coldest of all models in the year 2059, but nonetheless has a long-term trend that sits approximately in the middle of the ensemble.

Figure 5-27 should not be interpreted as a set of specific predictions for individual years. But it illustrates that although we expect a long term overall upward trend in temperatures (at least for RCP 8.5), there will still be some relatively cool years. However for this particular example, a year which is unusually warm under our present climate could become the norm by about 2050, and an “unusually warm” year in 30-50 years’ time (under the higher emission scenarios) is likely to be warmer than anything we currently experience.




Figure 5-27: New Zealand Temperature - historical record and an illustrative schematic projection illustrating future year-to-year variability. (See text for full explanation). After Mullan et al. (2016).

For rainfall, the fact that we have recently moved into a negative phase of the Interdecadal Pacific Oscillation (Figure 4-13) may be just as important for the Horizons Region over the next 2-3 decades as the effects of anthropogenic climate change. From Section 4.3.1, it can be seen that periods of negative SOI (e.g. El Niño) may on average experience slightly below normal rainfall in the eastern part of Horizons Region (east of Ruahine and Tararua Ranges), pushing rainfall in these directions in the same direction as expected from anthropogenic factors (Section 5.3). A subsequent further reversal of the IPO in 20-30 years' time could have the opposite effect, offsetting part of the anthropogenic trend in rainfall for a few decades.

As discussed in Section 4.3, the IPO and the El Niño/La Niña cycle have an effect on New Zealand sea level. So the sea levels we experience over the coming century will also result from the sum of anthropogenic trend and natural variability.

The message from the section is *not* that anthropogenic trends in climate can be ignored because of natural variability. In the projections we have discussed these anthropogenic trends become the dominant factor locally as the century progresses. Nevertheless, we need to bear in mind that at some times natural variability will be adding to the human-induced trends, while at others it may be offsetting part of the anthropogenic effect.

6 Horizons Region – Impacts, Vulnerability and Adaptation

The main purpose of this report has been to draw together existing information on how Horizons Region's climate may change in the future. The resourcing did not extend to undertaking a detailed evaluation of the likely impacts of these changes, of the vulnerability of the Horizons Region to these impacts, or of investigating options for adapting to them.

Ways in which councils can investigate some of the physical climate issues are outlined in the guidance manual published by the Ministry for the Environment (Ministry for the Environment, 2008a). These have not been updated in the recent report for the Ministry (Mullan et al., 2016) as the material is considered to be excellent guidance and still relevant to the new projections. The report on coastal hazards and climate change is also useful (Ministry for the Environment, 2008b) as is the erosion report by Basher et al. (2012).

The Ministry for the Environment climate change guidance manuals recommend that councils should build consideration of climate change into their planning activities rather than considering them in isolation, and should take a risk management approach. Issues surrounding climate change impacts, especially related to local government as well as Maori communities, are covered by Manning et al. (2014). As illustrated by Figure 6-1, consideration of climate change becomes particularly important for designing climate-sensitive infrastructure or assets which are likely to be around for many decades, and for resource use and land development planning over similar timescales.




Figure 6-1: Time scales and adaptation. Planning for human-induced climate change becomes increasingly important as one moves right along this line. Source: R. Jones, CSIRO.

Some particular impact, vulnerability, and adaptation issues to which Horizons Regional Council may wish to give consideration include:

- Implications of sea-level rise and coastal change for planning and development in coastal areas.
- Implications of river flow and sedimentation changes, as well as changing flood regimes for river engineering planning.
- Implications of potential changes in rainfall and of drought frequency for water demand, availability and allocation (including planning for irrigation schemes and storage).
- Implications of projected changes in extreme rainfall, erosion risk and coastal hazards for council roading and stormwater drainage infrastructure, lifelines planning, and civil defence and emergency management.
- Opportunities which climate change may bring for new horticultural crops – and infrastructure and land-use issues that might arise.
- Implications of climate change (including potential changes in flood frequency, extreme rainfall (influencing hillslope and riverbank erosion) and in coastal hazards) for land-use planning (e.g. erosion control measures, Basher et al. (2012), Manderson et al. (2015)).
- Implications for natural ecosystems and their management, both terrestrial and marine. This is especially relevant given the two National Parks and a number of forest parks in the region. Reisinger (2014) gives information on the projected impacts on natural ecosystems for New Zealand as a whole.
- Building consideration of climate change impacts and adaptation into council planning as outlined in MfE guidance. Also important is consultation and discussion with stakeholders (e.g. groups of farmers, iwi) to help them identify climate-related risks and ways of building resilience (e.g. King et al. (2013)).

Recommendations for future work that may assist Horizons Regional Council in understanding potential climate change-related impacts:

- Influence of climate-induced coastal hazard drivers such as sea-level rise, storm surge and wave climate on Horizon Region's coastlines.
- River flow projections for additional rivers in Horizons Region.

7 Acknowledgements

The following NIWA staff are acknowledged for their assistance in compiling this report: Andrew Tait, Nicolas Fauchereau, Aran Mol, Ben Robinson, and Scott Stephens.

8 References

- BASHER, L., ELLIOTT, S., HUGHES, A., TAIT, A., PAGE, M., ROSSER, B., MCIVOR, I., DOUGLAS, G. & JONES, H. 2012. Impacts of climate change on erosion and erosion control methods - A critical review. MPI Technical Paper No: 2012/45, 216pp.
- BATES, N. R., ASTOR, Y. M., CHURCH, M. J., CURRIE, K., DORE, J. E., GONZALEZ-DAVILA, M., LORENZONI, L., MULLER-KARGER, F., OLAFSSON, J. & SANTANA-CASLANO, J. M. 2014. A time-series view of changing ocean chemistry due to ocean uptake of anthropogenic CO₂ and ocean acidification. *Oceanography*, 27, 126-141.
- BELL, J. 2014. Coastal hazard assessment of Koitiata and Castlecliff Beaches: Rangitikei and Wanganui Districts, report for Horizons Regional Council, 2014/EXT/1379.
- BELL, J. 2015. Himatangi Beach Coastal Hazards Assessment, Manawatu District. Report for Horizons Regional Council, 2015/EXT/1422.
- BELL, J. J., DAVY, S. K. & JONES, T., ET AL. 2013. Could some coral reefs become sponge reefs as our climate changes? *Global Change Biology* 19, 2613-2624
- BLACKWOOD, P. 2007. Storm surge and wave run up design levels for Foxton Beach: An assessment of flood risks and mitigation options, report for Horizons Regional Council, 2007/INT/497.
- BOOKER, D. J. 2013. Spatial and temporal patterns in the frequency of events exceeding three times the median flow (FRE3) across New Zealand. *Journal of Hydrology (NZ)*, 52, 15-40.
- BOSTOCK, H. C., TRACEY, D. M., CURRIE, K. I., DUNBAR, G. B., HANDLER, M. R., MIKALOFF-FLETCHER, S. E., SMITH, A. M. & WILLIAMS, M. J. M. 2015. The carbonate mineralogy and distribution of habitat-forming deep-sea corals in the Southwest Pacific region. . *Deep-Sea Research I* 100, 88–104.
- BURRELL, T. J. 2015. *Bacterial extracellular Enzyme Activity in a Future Ocean*. PhD thesis, Victoria University of Wellington.
- BYLENGA, C., CUMMINGS, V. J. & RYAN, K. 2015. Fertilisation and larval development in the Antarctic bivalve, *Laternula elliptica* under reduced pH and elevated temperatures. *Marine Ecology Progress Series*, 536, 187-201.
- BYRNE, M., LAMARE, M., WINTER, D., DWORJANYN, S. A. & UTHICKE, S. 2013. The stunting effect of a high CO₂ ocean on calcification and development in sea urchin larvae, a synthesis from the tropics to the poles. *Phil Trans R Soc B* 368.
- CHAPPELL, P. R. 2012. The climate and weather of Manawatu-Wanganui. NIWA Science and Technology Series: 66, 40 pp.
- CHURCH, J. A., CLARK, P. U., CAZENAVE, A., GREGORY, J. M., JEVREJEVA, S., LEVERMANN, A., MERRIFIELD, M. A., MILNE, G. A., NEREM, R. S., NUNN, P. D., PAYNE, A. J., PFEFFER, W. T., STAMMER, D. & UNNIKRISHNAN, A. S. 2013a. Sea-level rise by 2100. *Science*, 342, 1445.
- CHURCH, J. A., CLARK, P. U., CAZENAVE, A., GREGORY, J. M., JEVREJEVA, S., LEVERMANN, A., MERRIFIELD, M. A., MILNE, G. A., NEREM, R. S., NUNN, P. D., PAYNE, A. J., PFEFFER, W. T., STAMMER, D. & UNNIKRISHNAN, A. S. 2013b. Sea Level Change. In: STOCKER, T. F., QIN, D., PLATTNER, G.-K., TIGNOR, M., ALLEN, S., BOSCHUNG, J., NAUELS, A., XIA, Y., BEX, V. & MIDGLEY, P. M. (eds.) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- CLARK, A., MULLAN, B. & PORTEOUS, A. 2011. Scenarios of Regional Drought under Climate Change. Wellington: NIWA Report WLG2010-32 for Ministry of Agriculture and Forestry.
- CLARK, D., LAMARE, M. & BARKER, M. 2009. Response of sea urchin pluteus larvae (Echinodermata: Echinoidea) to reduced seawater pH: a comparison among a tropical, temperate, and a polar species. *Marine Biology* 156, 1125-1137.

- CUMMINGS, V., HEWITT, J., ALLEN, S., MARRIOTT, P., BARR, N. & HEATH, P. 2013. Ocean acidification: impacts on key NZ molluscs, with a focus on flat oysters (*Ostrea chilensis*). Final Research Report for the NZ Ministry of Primary Industries on project ZBD200913. 18 p.
- CUMMINGS, V., HEWITT, J., VAN ROOYEN, A., CURRIE, K. & BEARD, S., ET AL. 2011. Ocean Acidification at High Latitudes: Potential Effects on Functioning of the Antarctic Bivalve *Laternula elliptica*. . *PLoS ONE*, 6, e16069.
- CUMMINGS, V., SMITH, A., PEEBLES, B. & MARRIOTT, P. 2015. Shell mineralogy of flat oysters under ocean acidification conditions: preliminary report. Progress report to Ministry of Primary Industries Project ZBD201306. June. 9 p.
- CUNNINGHAM, S. C. 2013. *The effects of ocean acidification on juvenile Haliotis iris*. Master of Science, University of Otago.
- DEAN, S., ROSIER, S., CAREY-SMITH, T. & STOTT, P. A. 2013. The role of climate change in the two-day extreme rainfall in Golden Bay, New Zealand, December 2011. *Bulletin of the American Meteorological Society*, 94, 561-563.
- GOODHUE, N. G., ROUSE, H. L., RAMSAY, D., BELL, R. G., HUME, R. M. & HICKS, D. M. 2012a. Coastal adaptation to climate change: mapping a New Zealand coastal sensitivity index. A report prepared as part of the Coastal Adaptation to Climate Change Project under contract (CO1XO802) to MBIE, NIWA, Hamilton, New Zealand. 43p.
- GOODHUE, N. G., STEPHENS, S. A. & GORMAN, R. G. 2012b. Combined wave and storm tide hazard for southern Tasman Bay. NIWA report HAM2012-087 prepared for Nelson City Council
- GRIFFITHS, G. M. 2006. Changes in New Zealand daily rainfall extremes. *Weather and Climate*, 27, 3-44.
- HANNAH, J. & BELL, R. G. 2012. Regional Sea Level Trends in New Zealand. *Journal of Geophysical Research: Oceans*, 117, 1-7.
- HENDERSON, R. & DIETTRICH, J. 2007. Statistical analysis of river flow data in the Horizons Region, NIWA Client Report for Horizons Regional Council, CHC2006-154, 349pp.
- HEPBURN, C. D., PRITCHARD, D. W. & CORNWALL, C. E., ET AL. 2011. Diversity of carbon use strategies in a kelp forest community: implications for a high CO₂ ocean. . *Global Change Biology* 17, 2488-2497.
- IPCC 2013. Summary for Policymakers. In: STOCKER, T. F., QIN, D., PLATTNER, G.-K., TIGNOR, M., ALLEN, S. K., BOSCHUNG, J., NAUELS, A., XIA, Y., BEX, V. & MIDGLEY, P. M. (eds.) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- IPCC 2014a. Summary for Policymakers. In: FIELD, C. B., BARROS, V. R., DOKKEN, D. J., MACH, K. J., MASTRANDREA, M. D., BILIR, T. E., CHATTERJEE, M., EBI, K. L., ESTRADA, Y. O., GENOVA, R. C., GIRMA, B., KISSEL, E. S., LEVY, A. N., MACCRACKEN, S., MASTRANDREA, P. R. & WHITE, L. L. (eds.) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York, NY, USA: Cambridge University Press.
- IPCC 2014b. Summary for Policymakers. In: EDENHOFER, O., R. PICHES-MADRUGA, Y. SOKONA, E. FARAHANI, S. KADNER, K. SEYBOTH, A. ADLER, I. BAUM, S. BRUNNER, P. EICKEMEIER, B. KRIEMANN, J. SAVOLAINEN, S. SCHLÖMER, C. VON STECHOW, T. ZWICKEL & MINX, J. C. (eds.) *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA.: Cambridge University Press.
- JAMES, R. K., HEPBURN, C. D. & CORNWALL, C. E., ET AL. 2014. Growth response of an early successional assemblage of coralline algae and benthic diatoms to ocean acidification.. *Marine Biology* 161, 1687-1696

- KIDSTON, J., RENWICK, J. A. & MCGREGOR, J. 2009. Hemispheric-scale seasonality of the Southern Annular Mode and impacts on the climate of New Zealand. *Journal of Climate*, 22, 4759-4770.
- KING, D. N., DALTON, W., BIND, J., SRINIVASAN, M. S., HICKS, D. M., ITI, W., SKIPPER, A., HOME, M. & ASHFORD-HOSKING, D. 2013. Coastal adaptation to climate vulnerability and change: examining community risk, vulnerability and endurance at Mitimiti, Hokianga, Aotearoa-New Zealand. NIWA report AKL2013-022.
- LAW, C. S., BREITBARTH, E., HOFFMANN, L. J., MCGRaw, C. M., LANGLOIS, R. J., LAROCHE, J., MARRINER, A. & SAFI, K. A. 2012. No stimulation of nitrogen fixation by non-filamentous diazotrophs under elevated CO₂ in the South Pacific. *Global Change Biology*, 18, 3004–3014.
- MAAS, E. W., HALL, J. A., LAW, C. S., PICKMERE, S., CURRIE, K. I., CHANG, F. H., VOYLES, K. M. & CAIRD, D. 2013. Effect of ocean acidification on bacterial abundance, activity and diversity in the Ross Sea, Antarctica. *Appl. Microbial Ecol.*, 70, 1-15.
- MANDERSON, A., DYMOND, J. & AUSSEIL, A.-G. 2015. Climate change impacts on water quality outcomes from the Sustainable Land Use Initiative (SLUI), Report for Horizons Regional Council by Landcare Research, 2015/EXT/1451, 39pp.
- MANNING, M., LAWRENCE, J., KING, D. N. & CHAPMAN, R. 2014. Dealing with changing risks: a New Zealand perspective on climate change adaptation. *Regional Environmental Change*, 15, 581-594.
- MINISTRY FOR THE ENVIRONMENT 2008a. Climate Change Effects and Impacts Assessment. A Guidance Manual for Local Government in New Zealand. In: MULLAN, B., WRATT, D., DEAN, S., HOLLIS, M., ALLAN, S., WILLIAMS, T., KENNY, G. & MFE (eds.). Wellington: Ministry for the Environment.
- MINISTRY FOR THE ENVIRONMENT 2008b. Coastal hazards and climate change. A guidance manual for local government in New Zealand. 2nd edition. Revised by Ramsay, D. and Bell, R. (NIWA). Prepared for Ministry for the Environment. viii+127 p.
- MULLAN, A. B., PORTEOUS, A., WRATT, D. & HOLLIS, M. 2005. Changes in drought risk with climate change. NIWA report WLG2005-23 for Ministry for the Environment and Ministry of Agriculture and Fisheries, Wellington.
- MULLAN, A. B., SOOD, A. & STUART, S. 2016. Climate Change Projections for New Zealand based on simulations undertaken for the IPCC 5th Assessment. Part A: Atmosphere Changes. NIWA Client Report for Ministry for the Environment, WLG2015-31. June 2016.
- MUNDAY, P. L., CHEAL, A. J. & DIXSON, D. L., ET AL. 2014. Behavioural impairment in reef fishes caused by ocean acidification at CO₂ seeps. *Nature Climate Change* 4, 487-492.
- REISINGER, A., KITCHING, R. L., CHIEW, F., HUGHES, L., NEWTON, P. C. D., SCHUSTER, S. S., TAIT, A. & WHETTON, P. 2014. Australasia. In: BARROS, V. R., FIELD, C. B., DOKKEN, D. J., MASTRANDREA, M. D., MACH, K. J., BILIR, T. E., CHATTERJEE, M., EBI, K. L., ESTRADA, Y. O., GENOVA, R. C., GIRMA, B., KISSEL, E. S., LEVY, A. N., MACCRACKEN, S., MASTRANDREA, P. R. & WHITE, L. L. (eds.) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK, and New York, NY, USA: Cambridge University Press.
- ROYAL SOCIETY OF NEW ZEALAND 2016. Climate change implications for New Zealand, <http://www.royalsociety.org.nz/expert-advice/papers/yr2016/climate-change-implications-for-new-zealand/>, 72p.
- SALINGER, M. J., RENWICK, J. A. & MULLAN, A. B. 2001. Interdecadal Pacific Oscillation and South Pacific climate. *International Journal of Climatology*, 21, 1705-1721.
- SHAND, R. 2008. Foxton Beach erosion hazard assessment, report for Horizons Regional Council by Coastal Systems Ltd., 2008-10 REP.

- SMITH, A. M., BERMAN, J. & KEY JR, M. M., ET AL. 2013. Not all sponges will thrive in a high-CO₂ ocean: Review of the mineralogy of calcifying sponges. *Palaeogeography Palaeoclimatology Palaeoecology*, 392, 463-472.
- TAIT, A., HENDERSON, R., TURNER, R. & ZHENG, X. G. 2006. Thin plate smoothing spline interpolation of daily rainfall for New Zealand using a climatological rainfall surface. *International Journal of Climatology*, 26, 2097-2115.
- TAIT, L. 2014. Impacts of natural and manipulated variations in temperature, pH and light on photosynthetic parameters of coralline–kelp assemblages. *J. Mar Exp Biol. Ecol.*, 454, 1-8.
- THOMPSON, D. W. J., SOLOMON, S., KUSHNER, P. J., ENGLAND, M. H., GRISE, K. M. & KAROLY, D. J. 2011. Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change. *Nature Geoscience*, 4, 741-749.
- THRESHER, R. E., GUINOTTE J.M., MATEAR R.J.; ET AL. 2015. Options for managing impacts of climate change on a deep-sea community. *Nature Climate Change* 5, 635-639.
- TITTENSOR, D. P., BACO, A. R. & HALL-SPENCER, J. M., ET AL. 2010. Seamounts as refugia from ocean acidification for cold-water stony corals. *Marine Ecology- An Evolutionary Perspective* 31, 212-225.
- TONKIN & TAYLOR 2010. Coastal Hazard Report: Akitio and Herbertville, report for Horizons Regional Council, 61457.
- TONKIN & TAYLOR 2013. Coastal Hazard Assessment: Waikawa to Waiterere, report for Horizons Regional Council, 61457.001.
- WONG, P. P., LOSADA, I. J., GATTUSO, J.-P., HINKEL, J., KHATTABI, A., MCINNES, K. L., SAITO, Y. & SALLINGER, A. 2014. Coastal systems and low-lying areas. In: BARROS, V. R., FIELD, C. B., DOKKEN, D. J., MASTRANDREA, M. D., MACH, K. J., BILIR, T. E., CHATTERJEE, M., EBI, K. L., ESTRADA, Y. O., GENOVA, R. C., GIRMA, B., KISSEL, E. S., LEVY, A. N., MACCRACKEN, S., MASTRANDREA, P. R. & WHITE, L. L. (eds.) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York, NY, USA: Cambridge University Press.